



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/47, A61K 38/17		A1	(11) International Publication Number: WO 00/23584																																																									
			(43) International Publication Date: 27 April 2000 (27.04.00)																																																									
(21) International Application Number: PCT/EP99/07832 (22) International Filing Date: 15 October 1999 (15.10.99) (30) Priority Data: 98119583.7 16 October 1998 (16.10.98) EP (71) Applicants (for all designated States except US): BOEHRINGER INGELHEIM INTERNATIONAL GMBH [DE/DE]; Postfach 200, D-55216 Ingelheim am Rhein (DE). UNIVERSITY HOSPITAL LEIDEN [NL/NL]; Rijnsburgerweg 10, NL-2321 RP Leiden (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): SCHRIER, Peter, I. [NL/NL]; Schelpenkade 17, NL-2313 ZT Leiden (NL). AARNOUDSE, Corlien, A. [NL/NL]; Pr. Alexanderlaan 18, NL-2224 XM Katwijk (NL). HEIDER, Karl-Heinz [DE/AT]; Johann-Strauss-Promenade 4/11, A-2000 Stock- erau (AT). KLADE, Christoph [AT/AT]; Gröhrmühlgasse 1b/17, A-2700 Wr. Neustadt (AT). (74) Agents: LAUDIEN, Dieter; Boehringer Ingelheim GmbH, D-55216 Ingelheim am Rhein (DE) et al.		(81) Designated States: CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.																																																										
(54) Title: CAMEL, AN ALTERNATIVE TRANSLATION PRODUCT OF THE TUMOUR ANTIGEN LAGE-I <table border="0"> <tr> <td>CAMEL</td> <td>-----</td> <td></td> </tr> <tr> <td>LAGE-1^S</td> <td>--ATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG</td> <td>48</td> </tr> <tr> <td>LAGE-1^L</td> <td>GCATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG</td> <td>50</td> </tr> <tr> <td>NY-ESO-1</td> <td>--ATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG</td> <td>48</td> </tr> <tr> <td>CAMEL</td> <td>-----CGACGGGCGATGCT</td> <td>14</td> </tr> <tr> <td>LAGE-1^S</td> <td>GAGCCATGCAGGCCGAAGGCCAGGGCACAGGGGGTTCGACGGGCGATGCT</td> <td>98</td> </tr> <tr> <td>LAGE-1^L</td> <td>GAGCCATGCAGGCCGAAGGCCAGGGCACAGGGGGTTCGACGGGCGATGCT</td> <td>100</td> </tr> <tr> <td>NY-ESO-1</td> <td>GAGCCATGCAGGCCGAAGGCCAGGGGGTTCGACGGGCGATGCT</td> <td>98</td> </tr> <tr> <td></td> <td>*****</td> <td></td> </tr> <tr> <td>CAMEL</td> <td>GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG</td> <td>64</td> </tr> <tr> <td>LAGE-1^S</td> <td>GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG</td> <td>148</td> </tr> <tr> <td>LAGE-1^L</td> <td>GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG</td> <td>150</td> </tr> <tr> <td>NY-ESO-1</td> <td>GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG</td> <td>148</td> </tr> <tr> <td></td> <td>*****</td> <td></td> </tr> <tr> <td>CAMEL</td> <td>CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG</td> <td>114</td> </tr> <tr> <td>LAGE-1^S</td> <td>CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG</td> <td>198</td> </tr> <tr> <td>LAGE-1^L</td> <td>CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG</td> <td>200</td> </tr> <tr> <td>LAGE-1^L</td> <td>CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG</td> <td>198</td> </tr> <tr> <td></td> <td>*****</td> <td></td> </tr> </table>				CAMEL	-----		LAGE-1 ^S	--ATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG	48	LAGE-1 ^L	GCATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG	50	NY-ESO-1	--ATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG	48	CAMEL	-----CGACGGGCGATGCT	14	LAGE-1 ^S	GAGCCATGCAGGCCGAAGGCCAGGGCACAGGGGGTTCGACGGGCGATGCT	98	LAGE-1 ^L	GAGCCATGCAGGCCGAAGGCCAGGGCACAGGGGGTTCGACGGGCGATGCT	100	NY-ESO-1	GAGCCATGCAGGCCGAAGGCCAGGGGGTTCGACGGGCGATGCT	98		*****		CAMEL	GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG	64	LAGE-1 ^S	GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG	148	LAGE-1 ^L	GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG	150	NY-ESO-1	GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG	148		*****		CAMEL	CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG	114	LAGE-1 ^S	CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG	198	LAGE-1 ^L	CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG	200	LAGE-1 ^L	CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG	198		*****	
CAMEL	-----																																																											
LAGE-1 ^S	--ATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG	48																																																										
LAGE-1 ^L	GCATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG	50																																																										
NY-ESO-1	--ATCCTCGTGGGCCCTGACCTTCTCTCTGAGAGCCGGGCAGAGGCTCCG	48																																																										
CAMEL	-----CGACGGGCGATGCT	14																																																										
LAGE-1 ^S	GAGCCATGCAGGCCGAAGGCCAGGGCACAGGGGGTTCGACGGGCGATGCT	98																																																										
LAGE-1 ^L	GAGCCATGCAGGCCGAAGGCCAGGGCACAGGGGGTTCGACGGGCGATGCT	100																																																										
NY-ESO-1	GAGCCATGCAGGCCGAAGGCCAGGGGGTTCGACGGGCGATGCT	98																																																										

CAMEL	GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG	64																																																										
LAGE-1 ^S	GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG	148																																																										
LAGE-1 ^L	GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG	150																																																										
NY-ESO-1	GATGGCCCAGGAGGCCCTGGCATTCTGATGGCCCAGGGGGCAATGCTGG	148																																																										

CAMEL	CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG	114																																																										
LAGE-1 ^S	CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG	198																																																										
LAGE-1 ^L	CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG	200																																																										
LAGE-1 ^L	CGGCCAGGAGAGGCGGGTGCCACGGGCGGCAGAGGTCCCCGGGGCGCAG	198																																																										

(57) Abstract <p>The tumor-associated antigen CAMEL and DNA encoding it. The tumor-associated antigen is encoded by an open reading frame of the LAGE-I gene. The tumor-associated antigen, immunogenic (poly)peptides derived therefrom and DNAs encoding them, are useful for cancer immunotherapy.</p>																																																												

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CAMEL, AN ALTERNATIVE TRANSLATION PRODUCT OF THE TUMOUR ANTIGEN LAGE-1

The present invention relates to the field of cancer therapy, more specifically to tumor-associated antigens.

Cytotoxic T lymphocytes (CTLs) play an important role in the defense
5 against melanoma. Melanoma-specific CTL clones have been obtained
from either tumor infiltrating lymphocytes (TIL) *in vitro* stimulated with
cytokines, or from peripheral blood mononuclear cells (PBMC) cultured with
(autologous) tumor cells. T cell responses against tumor cells are enhanced
by cytokine transfection of the tumor cells, both in animal models and in *in*
10 *vitro* human culture systems. (van Elsas et al., 1997; Gansbacher et al.,
1990; Tepper et al., 1989; Fearon et al., 1990; Dranoff et al., 1993)

The antigens recognized by the tumor-specific T cells become better
defined by the development of molecular cloning techniques. These T cell
targets can be divided in three groups: 1) tumor-specific antigens, not
15 expressed in healthy tissues, except testis and placenta (e.g., MAGE,
BAGE, GAGE, NY-ESO-1, LAGE-1); 2) antigens that are lineage-specific
and expressed in both melanoma and melanocytes (e.g., MART-1/ Melan-
A, gp100, tyrosinase) and 3) unique, mutated antigens (e.g., β -catenin,
CDK4, MUM-1) (reviewed by Van den Eynde and Brichard, 1995).

20 By means of Representational Difference Analysis (RDA), a PCR-based
method that has been used to identify genes with tissue-specific or tumor-
specific expression, the LAGE-1 and NY-ESO1 genes were identified as
being tumor specific by screening cDNA libraries from melanoma cell lines
with a primer from a cDNA clone enriched in melanoma-specific sequences
25 (Lethe et al., 1998).

NY-ESO-1 is a gene originally identified by SEREX technology (Chen et al.,
1997). It was shown to have two different reading frames (DNA sequences
and derived protein sequences given in SEQ ID NO: 7 - 10), translation

products of which were shown to contain epitopes of tumor specific T-cells (Jäger et al., 1998; Wang et al., 1998).

It was an object of the present invention to provide a novel tumor-associated antigen.

- 5 To solve the problem underlying the invention, melanoma cell line 518A2 and its IL-2- or GM-CSF-transfectants were compared for their CTL stimulating capacity *in vitro*. Stimulation of autologous PBMC with the IL-2 producing melanoma cells resulted in a melanoma-specific CTL response (van Elsas et al., 1997). CTL clones derived from this culture recognized,
10 besides autologous melanoma cell lines, also a panel of HLA-A*0201 positive melanoma cell lines, but were not reactive with normal melanocytes. Although 518A2 was shown to express a number of antigens previously identified to be recognized by anti-melanoma CTL (van Elsas et al., 1996), the CTL clones available recognize a new melanoma-specific
15 antigen that is immunodominant in 518A2.

- In the experiments of the present invention, the target structure that was recognized by one of these CTL clones was fully characterized and named CAMEL (CTL-recognized Antigen on Melanoma). These sequences are described in the attached sequence listing as SEQ ID NO: 1 and SEQ ID
20 NO: 2.

It was surprisingly found that CAMEL is encoded by a reading frame of the LAGE-1^s-cDNA (SEQ ID NO:3) that is distinct from that encoding the putative LAGE-1 protein (SEQ ID NO: 4). (This reading frame is designated ORF-1.)

- 25 In the present invention, a cDNA clone was identified that lacks the first 84 bp of the LAGE-1^s sequence (SEQ ID NO: 3) which means that it is devoid of the initiation codon at position 54 of that sequence (Fig. 2a). The first possible translation initiation site in this clone (4H8) is the ATG at position 94 of LAGE-1^s (SEQ ID NO: 3), which is however, not in frame with

the first ATG at position 54. Therefore, the CAMEL protein (SEQ ID NO: 2) translated from the 4H8 cDNA clone is different from the putative LAGE-1^S protein (SEQ ID NO: 4).

In a first aspect, the present invention is directed to the tumor-associated antigen CAMEL (SEQ ID NO: 2) which is encoded by an isolated DNA molecule with the sequence as defined in SEQ ID NO: 1.

The coding sequence of CAMEL corresponds to the ORF-1 of LAGE-1 cDNA (Lethe et al., 1997; WO 98/32855).

In the present invention „ORF-1“ is defined as the open reading frame starting with ATG at position 94 of SEQ ID NO:3 (LAGE-1^S), which corresponds to position 10 in SEQ ID NO: 1 (CAMEL), to position 96 in SEQ ID NO: 5 (LAGE-1^L)

In a further aspect, the present invention relates to immunogenic (poly)peptides derived from CAMEL. A first group of peptides is selected from peptides inducing a humoral immune response (induction of antibodies). Such peptides are selected by randomly choosing continuous stretches of amino acids (at least 12-15), applying them to an individual and confirming the generation of antibodies by standard immunological assays, e.g. ELISA. This group of immunogenic (poly)peptides also encompasses the entire CAMEL antigen or larger fragments thereof.

The second group of peptides, which is preferred, can be presented by MHC molecules (in humans: HLA molecules), they have the potential to induce an immune response, in particular by eliciting a CTL response.

In a preferred embodiment, immunogenic peptides which have the ability to elicit a CTL response, are selected from peptides with the sequence set forth in SEQ ID NO: 11, 12, 24, 25 and 26.

To obtain peptides that have the ability to elicit a cellular immune response, the selection of peptide sequences from a given antigen is, in the first place, based on the requirement for such peptide to bind to an MHC molecule present in the repertoire of the patient to be treated. Two classes of MHC
5 molecules are distinguished, class I and class II. Class I molecules consist of a membrane-inserted heavy chain and a non-covalently attached light chain. In their structure, MHC class I molecules resemble a moose's head, the antlers forming a groove which is recognized by the peptide. In humans, HLA-A, B and C are the "classical" MHC class I molecules.

10 Additional immunogenic peptides may be identified by methods known in the art which rely on the correlation between MHC-binding and CTL induction, e.g. those used by Stauss et al., 1992, who identified candidate T-cell epitopes in human papilloma virus.

Since immunogenic peptides can be predicted based on their "peptide
15 binding motif" synthetic peptides which represent CTL epitopes may be designed and synthesized. Several methods, which are useful in the present invention for designing peptides, have been used to identify CTL epitopes from known protein antigens.

It is well established that every MHC class I allelic product has
20 allele-specific requirements for the peptide ligand that binds to its groove and that it ultimately presents. These requirements were summarized as a motif by Falk et al., 1991. A large number of MHC peptide motifs and MHC ligands have become known to date. A method to search a known protein sequence for epitopes fitting to a given class I molecule, which is
25 based on this knowledge and which may be used in the present invention, was reviewed by Rammensee et al., 1995. It comprises the following steps: first, the protein sequence is screened for stretches fitting to the basic anchor motif (two anchors in most cases), whereby allowance should be made for some variation in peptide lengths as well as in anchor occupancy.
30 If a motif, for example calls for 9mers with Ile or Leu at the end, 10mers with

a fitting C-terminus should be considered as well, and other aliphatic residues at the C-terminus, like Val or Met, should also be considered. In this way, a list of peptide candidates is obtained. These are inspected for having as many non-anchor residues as possible in common with ligand
5 already known, or with the residues listed among the "preferred residues" or "others" on top of each motif (Table, given by Rammensee et al., 1995), for various HLA molecules. Binding assays can be performed at this stage to exclude weak binders which occur frequently among peptides conforming to a basic motif. If a detailed study on peptide binding requirements is
10 available, the candidates can also be screened for non-anchor residues detrimental or optimal for binding (Ruppert et al., 1993). One should keep in mind, however, that pure peptide binding motifs can be misleading in the search for natural ligands, since other constraints, such as enzyme specificity during antigen processing and specificity of transporters or
15 chaperones, are likely to contribute to ligand identity in addition to the MHC binding specificity.

This approach was successfully applied by, inter alia, Kawakami et al., 1995, to identify gp100 epitopes based on known HLA-A2.1 motifs. The validity of the method was confirmed by identifying, in parallel, the epitope
20 regions by using COS cells transfected with cDNA fragments generated by sequential deletion and testing for T-cell reactivity, as described above.

Recently, data bases and prediction algorithms have become available that enable to predict, with great reliability, peptide epitopes that bind to HLA molecules of interest.

25 Examples for peptide candidates with potential immunogenicity that can be derived from the tumor-associated antigen of the present invention, are the CAMEL-derived peptides with the sequence HLSPDQGRF and LMAQEALAF for HLA-A3 or RMAVPLLR for HLA-A3101. Similarly, other peptides for these or for other alleles can be determined by the method
30 mentioned above.

- The peptide binding can be tested in peptide binding assays. In order to determine the immunogenicity of the selected peptide or peptide equivalent, as defined below, which is the crucial parameter for peptide-based vaccine development and which in most cases strongly correlates with the stability of the peptide-MHC interaction (van der Burg et al., 1996), the methods described by Sette et al., 1994, in combination with quantitative HLA-binding assays, may be used. Alternatively, immunogenicity of the selected peptide may be checked by performing *in vitro* CTL induction by known methods e.g. as described below for *ex vivo* CTL induction.
- Alternatively to peptides derived from the naturally expressed tumor antigens, functional equivalents thereof, i.e. peptides with partially altered sequences or substances mimicking peptides, e.g. "peptidomimetics" or retro-inverso peptides, may be obtained by the following methods:
- To enhance the immunogenicity of the peptides, amino acid substitutions may be introduced at anchor positions to increase peptide MHC class I-binding affinity. The modified peptides are subsequently evaluated for enhanced binding and immunogenicity by screening for recognition by TIL (tumor-infiltrating lymphocytes) and CTL induction as described by Parkhurst et al., 1996, and Bakker et al., 1997.
- Another method useful in the present invention to find more immunogenic peptides by screening peptide libraries with a known CTL was described by Blake et al. 1996; it suggests the use of combinatorial peptide libraries for constructing functional mimics of tumor epitopes recognized by MHC class I-restricted CTLs.
- In principle, the selection of peptides capable of eliciting a cellular immune response is carried out in several steps, as described in WO 97/30721, which disclosure is incorporated herein by reference. In short, the candidates are first tested for their binding ability to an MHC molecule; subsequently good binders are tested for immunogenicity. A general

strategy for obtaining efficient immunogenic peptides has been described by Schweighoffer, 1997.

The polypeptide of the present invention or immunogenic peptides derived from its sequence, respectively, can be produced recombinantly or by peptide synthesis, as described in WO 96/10413, the disclosure of which is incorporated herein by reference. For recombinant production, a DNA molecule encoding the antigen or the CTL peptide of interest is inserted into an expression vector, transformed or transfected into an appropriate host cell, cultivated under conditions suitable for expression, recovered and purified. For chemical synthesis, various conventional techniques may be used, e.g. commercially available automatic synthesizers.

The tumor antigen of the present invention and the immunogenic peptides derived therefrom or the respective peptide equivalents are useful in cancer therapy, e.g. to induce, in the context of the appropriate MHC presenting molecule, an immunological response to tumors which express the corresponding antigen determinants. The induction of CTLs can be accomplished *in vivo* or *ex vivo*.

For *in vivo* induction of CTLs, a pharmaceutical composition comprising the peptide/antigen is administered to an individual suffering from a tumor associated with the respective tumor antigen in an amount sufficient to elicit an effective CTL response to the antigen-bearing tumor. Thus, the present invention provides pharmaceutical compositions for therapeutic treatment which are intended for parenteral, topical, oral or local administration. Preferably, the compositions are for parenteral administration, e.g. for intravenous, subcutaneous, intradermal or intramuscular application. The peptides/antigens are dissolved or suspended in a pharmaceutically acceptable carrier, preferably an aqueous carrier. The composition may contain additional auxiliary substances, e.g. buffering agents, etc. The peptides may be used alone or in combination with adjuvants, e.g. saponins, alum, or, in a particularly preferred embodiment, polycations,

like polyarginine or polylysine. The peptides may also be linked to components assisting CTL priming, e.g. T helper peptides, lipids or liposomes or coadministered with such components or with immunostimulating substances, e.g. cytokines (IL-2, IFN- γ). Methods and compositions for preparing and administering pharmaceutical compositions for therapeutic treatments are described in WO 95/04542 and WO 97/30721 the disclosures of which are herein incorporated by reference.

The immunogenic peptides may also be used to elicit a CTL response *ex vivo*. An *ex vivo* CTL response to a tumor expressing the antigen is induced by incubating a patient's CTL precursor cells together with antigen presenting cells and the immunogenic peptide. The thus activated CTLs are allowed to mature and expand to effector CTLs which are then readministered to the patient. Alternatively, the tumor antigen may be pulsed onto APCs which present MHC class II-reactive peptides (Mayordomo et al., 1995; Zitvogel et al., 1996). A suitable method for loading peptides onto cells, e.g. dendritic cells, is disclosed in WO 97/19169.

The peptides of the invention are preferably applied as a combination of peptides, e.g. different CAMEL-peptides. In an even more preferred embodiment, the peptides of the invention are combined with peptides derived from other tumor antigens, e.g. LAGE-1 and ESO-NY-1. The selection of the peptides is optimized towards covering multiple HLA types in order to be useful for a broad population of patients and/or towards a broad variety of malignancies, which is taken into account by combining peptides from a large variety of tumor antigens. The number of peptides suitable to be combined to yield an efficient therapy may vary within a broad range, e.g. from about 2 to approximately 100.

In a further aspect, the present invention is directed to an isolated DNA molecule with the sequence set forth in SEQ ID NO: 1 encoding CAMEL.

This DNA molecule, which is designated "CAMEL-DNA", contains the ORF-1 of LAGE-1 cDNA which is defined by nucleotides 54 - 336 of the sequence set forth in SEQ ID NO: 3.

The CAMEL-DNA of the present invention, or the corresponding RNA, may
5 be used, as an alternative to the use of the protein or the peptide, for cancer immunotherapy. Alternatively to using the natural sequence or fragments thereof, engineered derivatives may be utilized. These include sequences modified to encode (poly)peptides with improved immunogenicity, e.g. taking into account the modifications described above
10 for the peptides. Another form of modification is the assembly of multiple sequences encoding immunologically relevant peptides in a so-called string-of-beads fashion, as described by Toes et al., 1997. The sequences may also be modified by adding auxiliary coding elements, e.g. targeting functions that ensure more efficient delivery and processing of the
15 immunogen (e.g. Wu et al., 1995).

The nucleic acid molecules may be delivered either directly or as part of a recombinant virus or bacterium. In principle, any method that is known for gene therapy may be applied for nucleic acid-based cancer immunotherapy, both *in vivo* and *ex vivo*.

20 Examples for *in vivo* delivery are direct injection (injection of "naked" DNA) either intramuscularly or by "gene gun", which has been shown to result in the generation of CTLs against tumor antigens. Examples for recombinant organisms are vaccinia virus, fowlpox virus and adenovirus or *Listeria monocytogenes* (see Coulie, 1997 for a comprehensive review).
25 Furthermore, synthetic nucleic acid carriers like cationic lipids, microspheres, microbeads, liposomes may be useful for *in vivo* delivery of the sequence encoding respective antigen/peptide. Similarly as for peptides, various auxiliary agents that enhance the immune response may be co-applied, e.g. cytokines, either as proteins or as plasmids encoding
30 these.

Examples for *ex vivo* delivery are transfection of dendritic cells (Tuting, T., 1997) or other antigen presenting cells which are applied as a cellular cancer vaccine.

5 The present invention is also directed to the use of cells that express the tumor-associated antigen of the invention, either naturally or upon transfection with the respective coding sequence, for the preparation of a tumor vaccine.

10 In the present invention, it has been shown that CTL clones raised against IL-2 producing melanoma cell line 518/IL-2.14 are reactive against two alternatively spliced variants of LAGE-1, LAGE-1^S (SEQ ID NO: 3) and LAGE-1^L (SEQ ID NO: 5) and NY-ESO-1 (SEQ ID NO: 9). NY-ESO-1 is a recently described tumor antigen, identified by screening a cDNA library of an esophagus carcinoma with autologous patient serum (SEREX-method (Chen et al., 1997)). NY-ESO-1 is expressed in different tumor types but
15 not in healthy tissues except the testis.

In the present invention, the epitope of specific CTL 1/29 was determined by cDNA expression cloning and a truncated LAGE-1 cDNA clone was found. This truncation led to the identification of the peptide epitope in an alternative reading frame, since the "normal" translation initiation site of
20 LAGE-1 was absent. However, COS/HLA-A*0201 cells transfected with full length LAGE-1 or NY-ESO-1 cDNA clones could stimulate the CTL clone to TNF- α production as well. This probably means that two different proteins can be translated from one single mRNA.

25 NY-ESO-1 also has been described as the target of melanoma-specific HLA-A*0201 restricted CTL clones, which recognize an epitope translated in ORF3, located between aa 155 and 167 (Jäger et al., 1998). Therefore, it is very likely that also LAGE-1^S will be recognized by these clones, but not LAGE-1^L, since the protein sequence is different at this part of the molecule. The CAMEL-specific CTL clones recognize a peptide in an

alternative reading frame, which is encoded in both LAGE-1 and NY-ESO-1. This means that tumor cells expressing either LAGE-1 or NY-ESO-1 can be recognized by MLMAQEALAFI-specific CTL, which might enlarge the number of tumors that can be treated with immunotherapy based on this peptide.

Brief description of the Figures:

Figure 1: COS-7 transfection experiments with cDNA clone CAMEL and deletion constructs

- a) COS-7 cells were transfected with cDNAs as indicated and tested with CTL 1/29 in a TNF- α release assay.
- b) Deletion constructs of CAMEL cDNA were cotransfected with HLA-A*0201 cDNA in COS-7, followed by a TNF- α release assay with CTL 1/29. The PCR clones contain the numbers of nucleotides of the CAMEL cDNA as indicated.

Figure 2:

- a) Nucleotide alignment of cDNA clones CAMEL, LAGE-1^S, LAGE-1^L and NY-ESO-1.
- b) Protein translations of the cDNA clones LAGE-1^S, LAGE-1^L and NY-ESO-1. The translation of CAMEL is identical to the translation of LAGE-1^{S/L} in ORF1. Although ORF3 seems the most putative one, the CTL epitope is encoded in ORF1 (underlined).

Figure 3: Characterisation of peptides recognized by CTL clone 1/29

- a) TNF- α release assay with predicted HLA-A*0201 binding CAMEL peptides. Peptides as indicated were loaded on BLM, an

HLA-A*0201⁺ melanoma cell line, at a concentration of 10 µg/ml and tested with CTL 1/29 in a TNF-α release assay.

- b) The effects of increasing concentrations of peptides, derived from the major target epitope MLMAQEALAFI on recognition by CTL 1/29.

5 Various concentrations of peptides as indicated were loaded on HLA-A*0201⁺ cells and tested in a TNF-α release assay with CTL 1/29.

Figure 4: LAGE-1^{S/L} and NY-ESO-1 encode the CTL epitope

10 COS/HLA-A*0201 cells were transfected with these cDNA clones and reactivity with CTL 1/29 was measured in a TNF-α release assay.

Figure 5: His-tagged CAMEL protein, synthesized in E.coli

Figure 6: Expression of LAGE-1^{S/L} and NY-ESO-1 in healthy human tissues and melanoma cell lines

15 a) Northern Blot analysis of the expression of LAGE-1/NY-ESO-1 in a panel of healthy human tissues as indicated. The Blot was hybridised with ³²P-dCTP-labeled LAGE-1^S cDNA.

20 b) RT-PCR for LAGE-1 and NY-ESO-1. To discriminate between LAGE-1 and NY-ESO-1 mRNA, the same panel of melanoma cell lines was analysed by RT-PCR with gene-specific primers. Melanoma cell lines as indicated were used as targets in a TNF-α release assay with CTL 1/29.

Figure 7: Immunohistochemical analysis of CAMEL expression in human tumors

25 Figure 8: Stabilization of HLA-A2 surface expression by synthetic peptides on T2-cells

Brief description of the sequences:

	SEQ ID NO: 1:	CAMEL (4H8) cDNA sequence and translation
	SEQ ID NO: 2:	CAMEL protein sequence
5	SEQ ID NO: 3:	LAGE-1 ^S cDNA sequence and translation
	SEQ ID NO: 4:	LAGE-1 ^S protein sequence
	SEQ ID NO: 5:	LAGE-1 ^L cDNA sequence and translation
	SEQ ID NO: 6:	LAGE-1 ^L protein sequence
	SEQ ID NO: 7:	NY-ESO-1 cDNA sequence and translation
10	SEQ ID NO: 8:	NY-ESO-1 protein sequence
	SEQ ID NO: 9:	NY-ESO-1 cDNA and alternative translation
	SEQ ID NO: 10:	protein sequence of alternatively translated NY-ESO-1
	SEQ ID NO: 11:	peptide sequence of the CAMEL CTL epitope (11-mer)
	SEQ ID NO: 12:	peptide sequence of the CAMEL CTL epitope (10-mer)
15	SEQ ID NO: 13:	oligonucleotide SP6F-pSV
	SEQ ID NO: 14:	oligonucleotide R1
	SEQ ID NO: 15:	oligonucleotide R2
	SEQ ID NO: 16:	oligonucleotide T7R-pSV
	SEQ ID NO: 17:	oligonucleotide F3
20	SEQ ID NO: 18:	oligonucleotide ESO-1B
	SEQ ID NO: 19:	oligonucleotide ESO-1A
	SEQ ID NO: 20:	oligonucleotide 4H8-A
	SEQ ID NO: 21:	oligonucleotide 4H8-C
	SEQ ID NO: 22:	oligonucleotide CAMEL-XHO
25	SEQ ID NO: 23:	oligonucleotide CAMEL-KPN

- SEQ ID NO: 24 peptide CAMEL10
SEQ ID NO: 25 peptide CAMEL16
SEQ ID NO: 26 peptide CAMEL17
SEQ ID NO: 27 tyrosinase peptide
5 SEQ ID NO: 28 MAGE-3 peptide

In the Examples, if not stated otherwise, the following materials and methods were used

a) Cell cultures

- 10 Melanoma cell lines and COS-7 cells were maintained in DMEM containing 4.5 mM glucose supplemented with 8% FCS, 2 mM L-glutamine, 100 µg/ml of each penicillin and streptomycin. Melanoma cell line 518A2 was established from the dissected metastasis of a male patient in 1985, as described before (Versteeg et al., 1988). An IL-2 producing variant, 518/IL-
15 2.14, was obtained by transfection of 518A2 with the IL-2 cDNA (Osanto et al., 1993). Other melanoma cells that were used as targets in TNF- α release assay are FM3.29, FM6, FM28.4 and FM55P (gifts from J. Zeuthen, Denmark), MM127, MM415, MM485 (gifts from N. Hayward, Australia), SK-MEL-23, SK-MEL-29 (obtained from T. Wölfel, Mainz), Mi10221, Mi3046/2,
20 NA8, BLM (obtained from M. Visseren, Leiden). EBV-transformed B-LCL and the TNF- α -sensitive WEHI-164 clone 13 (a gift from Dr. P. Coulie, Brussels) were cultured in RPMI-1640, supplemented with L-glutamine and antibiotics as above, and 10% FCS.

- With the IL-2 producing cell line 518/IL-2.14 and autologous peripheral
25 blood mononuclear cells (PBMC) a CTL induction was performed, resulting in melanoma-specific HLA-A*0201 restricted CTL clones (van Elsas et al., 1997). The identification of the epitope of one of these clones, CTL 1/29, is reported here.

b) cDNA expression cloning

A cDNA library of 518/IL-2.14 was constructed in the expression vector pSVsport1 (GIBCO, BRL) using the Superscript Plasmid System (GIBCO, BRL). As to that purpose, poly-A⁺ mRNA was isolated using the Fast-Track system (Invitrogen), followed by reverse-transcription with an oligo-dT/NotI primer. Sall adapters were ligated to ds-cDNA and after NotI digestion and size fractionation, cDNA fragments were cloned into the pSVsport1 vector digested with Sall and NotI. After electroporation into ElectroMAX-DH10B (GIBCO, BRL) (following the manufacturers instructions) and selection for ampicillin resistance, 50-100 colonies were pooled for mini DNA isolation (QIAprep 8 plasmid kit, Qiagen). The in this way obtained cDNA pools were transfected in duplicate into COS-7 cells, together with the restriction element HLA-A*0201 (pBJ1.neo/HLA-A*0201, (Lin et al., 1990)), using the DEAE-dextran method. Briefly, COS-7 cells were seeded in 96-wells flatbottom plates at 1.5×10^4 cells per well in 100 μ l DMEM, 8% FCS. After 2 hours, medium was replaced by 30 μ l transfection mixture, containing 100 ng cDNA pool, 100 ng HLA-A*0201 cDNA, 400 ng/ml DEAE-dextran and 100 μ M chloroquine in serum free DMEM. Cells were incubated for 4 hours at 37°C and shocked for 2 minutes by the addition of 50 μ l 10% DMSO in PBS. The shock medium was replaced by 200 μ l DMEM, 8% FCS, and 48 hours later the cells were used as target cells for CTL in a TNF- α release assay.

c) Deletion constructs

Deletion constructs of cDNA clone 4H8 were obtained by PCR. PCR products were cloned in vector pCR3.uni (TA cloning system, Invitrogen). The constructs pCR-246 and pCR-464 were made with the vector-based forward primer, SP6F-pSV (SEQ ID NO: 13) and the reverse primers in cDNA 4H8, R1 (SEQ ID NO: 14) and R2 (SEQ ID NO: 15) respectively. As a control the complete 679 bp cDNA was cloned by PCR with two primers

on the pSVsport vector, SP6F-pSV (SEQ ID NO: 13) and T7R-pSV (SEQ ID NO: 16), resulting in pCR-679.

d) TNF- α release assay

CTL reactivity against tumor target cells, transfected COS-7 or peptide
5 loaded cells was measured in a TNF- α release assay. Target cells were seeded in duplicate or triplicate at $1.5\text{-}2 \times 10^4$ cells per well in a 96-wells flat bottom plate and 1500-2000 CTL were added to each well, in a total volume of 100 μl / well (IMDM, supplemented with antibiotics and 5% FCS). After 24 hours of co-culturing of effector and target cells, 50 μl out of each
10 well was added to a fresh 96-wells flatbottom plate, containing 50 μl (4.5×10^4) TNF- α -sensitive WEHI-164 cells per well in IMDM, supplemented with antibiotics, 5% FCS, 2 $\mu\text{g/ml}$ Actinomycin D and 40 mM LiCl. A viability staining was performed 24 hours later by the addition of 50 μl of
15 3-(4,5dimethylthiazol-2-yl)-2,5 diphenyltetrazolium bromide (MTT) solution (2.5 mg/ml in PBS). After incubation for 2-4 hours at 37°C the OD₅₅₀₋₆₅₀ was measured. TNF- α release in pg/ml was calculated from a standard with known TNF- α concentrations.

e) Northern Blot analysis

To determine expression in healthy tissues a Multiple Tissue Northern Blot
20 was obtained commercially (Clontech). As a probe, LAGE-1 cDNA was used, labeled with $\gamma\text{-}^{32}\text{P}$ -dCTP by use of the Mega-Prime Labeling kit (Amersham).

f) RT-PCR

cDNA synthesis was performed using oligo-dT and M-MLV reverse
25 transcriptase (Promega). Primers used for LAGE-1 specific PCR were the F3 (SEQ ID NO: 17) and ESO-1B primer (SEQ ID NO: 18). ESO-1B was also used as a reverse primer in the NY-ESO-1-specific PCR, while ESO-1A (SEQ ID NO: 19) was the forward primer in this reaction (Chen et

al., 1997). Reactions were performed in a Biometra-Uno or -Trio programmed as follows: 5 minutes 95°C, 30 cycles of 1 min. 95°C, 1 min. 58°C, 1 min. 72°C, followed by 10 minutes 72°C.

g) Expression of CAMEL in E. Coli

- 5 A fragment containing the coding sequence of CAMEL was made by PCR with the following primers:

4H8-A: GAAGAACATATGCTGATGGCCCAGGAGGC (SEQ ID NO: 20)

4H8-C: TTAAAGATCTCAGAACCGCCCCTGGTCG (SEQ ID NO: 21)

- This fragment was digested with NdeI and BglII and cloned in the NdeI and
10 BamHI sites of vector pET19b (Novagen, Madison, WI). This vector contains a 6xHis-tag coding sequence, allowing detection of the His-tagged protein with an anti-His antibody. The pET19b-CAMEL construct was transformed into BL21(DE3)pLysS E. coli bacteria (Novagen, Madison, WI). After culturing the bacteria at 30°C until an OD₆₀₀ = 0.5, IPTG (1 mM) was
15 added to induce overexpression of the His-tagged CAMEL protein. Samples were taken at 0h and 4h after IPTG induction and lysates of these samples were tested on a Western Blot with the Penta-His Antibody (Qiagen) according to the Western and Colony Blot protocol of the supplier. The His-tagged protein was visualized using the SuperSignal Substrate
20 system for Western blotting (Pierce, Rockford, US).

h) Preparation of anti-CAMEL antisera

Antibodies against the CAMEL protein were raised by immunizing two rabbits with three synthetic peptides derived from hydrophobic regions of this molecule :

- 25 F4: (K)GAMLAAQERRVPRAAEV(K) (pos. 15-31 of SEQ ID NO: 2)
A5: (K)GQQGPRGREEAPRGVRL(K) (pos. 36-52 of SEQ ID NO: 2)
B5: (K)KRRMEGAPAGPGGRRTAA(K) (pos. 58-73 of SEQ ID NO: 2)

(The lysine residues at both termini enable the peptides to be linked to KLH.)

For rabbit no. 1, 1 mg of each peptide was linked chemically to 2.5 mg of the carrier molecule KLH (Keyhole Limpet Hemocyanin) and after dialyzing, 0.8 mg of this protein in CFA (Complete Freund's Adjuvants) was weekly injected subcutaneously. Another rabbit (no. 2) was injected six times with the three peptides, not linked to KLH, following the scheme of 300 µg s.c. in CFA, 300 µg s.c. in IFA, 4x boost of 150 µg. The reactivity and specificity of the antisera were confirmed in ELISA and Western blot experiments. After four immunizations, antisera of both rabbits were reactive with the recombinant CAMEL protein synthesized in *E.coli*, but differed in their precise epitope: rabbit no. 1 produced antibodies against the CAMEL-B5 peptide, whereas the serum of rabbit no. 2 reacted with peptide F4. The antisera will further be referred to as "antiserum B5" and "antiserum F4".

j) Preparation of CAMEL-EGFP fusion proteins

The CAMEL coding sequence was fused to the *Aequorea victoria* -derived Green Fluorescent Protein (GFP). The CAMEL cDNA molecule was cloned into the pEGFP-N1 vector (Clontech), which contains a cDNA encoding the Enhanced, red shifted variant of GFP. In order to clone the cDNA molecule in frame with the EGFP cDNA and unidirectional, two primers were designed. The forward primer designated CAMEL-XHO (TTACTCGAGATGCTGATGGCCCAGG; SEQ ID NO: 22) covers the initiation codon ATG and contains an Xho1 site and the reverse primer CAMEL-KPN (AAGGTACCTTGAACCGCCCCTGGTCG; SEQ ID NO: 23) contains a mutation of the stop-codon and a Kpn1 site. The vector carrying the fusion construct was transfected into COS cells by calcium phosphate precipitation, protein lysates of the cells were used for Western blotting using CAMEL antisera against the CAMEL peptides B5 and F4, and anti-EGFP antibodies to detect the fusion protein according to standard protocols.

Example 1

cDNA clone 4H8 (CAMEL) encodes the target for melanoma-specific CTL 1/29

The antigenic epitope of melanoma-specific CTL 1/29 was identified by the
5 expression of cDNA library 518/IL2.14 and the restriction element HLA-A*0201 in COS-7 cells, followed by CTL screening in a TNF- α release assay. A positive pool of cDNAs was subcloned and clone 4H8, called CAMEL (SEQ ID NO: 1), was found to stimulate TNF- α release by the CTL to a similar extent as the original 518/IL2.14 cell line (Fig. 1). COS-7 cells or
10 COS-7 cells transfected with HLA-A*0201 or the 4H8 cDNA only were not recognized. The isolated 4H8 cDNA clone has a 679 bp insert, which shows strong homology with NY-ESO-1 (SEQ ID NO: 7), a tumor antigen originally identified as a target for humoral immune responses by serum screening methods (SEREX) (Chen et al., 1997). Colony hybridization of
15 the cDNA library, using clone 4H8 as a probe resulted in the detection of 2 types of full length clones which were called LAGE-1^S (SEQ ID NO: 3) and LAGE-1^L (SEQ ID NO: 5) (Fig. 2a). LAGE-1^L contains a 229 bp insertion at position 457, which has the consensus sequences for an intron, starting with a 5' GT and ending 3' AG. This indicates alternative splicing of
20 LAGE-1 mRNA. However, cDNA clone 4H8 lacks the first 84 bp of the LAGE-1 cDNA sequence.

Example 2

The peptide epitope of CTL 1/29 is coded in an alternative reading frame of LAGE-1 or NY-ESO-1

25 To identify which peptide was recognized by CTL 1/29, deletion constructs of cDNA 4H8 were transfected in HLA-A*0201⁺ COS-7 cells and tested in a TNF- α release assay. CTL reactivity was measured with all constructs (Fig. 1b), indicating that the epitope was coded within the first 330 bp of

clone 4H8. An HLA-A*0201 binding motif search was performed on the predicted protein sequence of that region (Drijfhout et al., 1995; D'Amaro et al., 1995), presuming that the ATG at position 10 in 4H8 functions as the translation initiation site. Predicted strong binding peptides at regions 1-11,
5 2-11, 1-9, 10-18, 11-19, 16-25, 17-25, 49-57, 55-63 and 70-78 of the CAMEL protein sequence (Fig. 2b) were added to HLA-A*0201⁺ BLM melanoma cells, and tested for CTL reactivity in a TNF- α release assay (Fig. 3b).

At a peptide concentration of 10 μ g/ml only the N-terminal 11- and 10-mer
10 peptides (M)LMAQEALAF_L (SEQ ID NO: 11 and NO: 12) induced preponderant recognition by CTL 1/29 (Fig. 3a), indicating that the epitope recognized by the CTL is located in the first 11 amino acids of the CAMEL-encoded protein. Closer inspection of peptides derived of this N-terminal 11-mer in a peptide concentration dependent TNF- α release assay (Fig. 3b)
15 revealed that the methionine at position 1 as well as the leucine at position 11 are of crucial importance for reconstituting CTL reactivity. Deletion of either of these amino acids leads to an at least 5 decades higher peptide concentration required for comparable TNF- α release. The only other peptide showing weak activity is the 3-11 MAQEALAF_L. In contrast, the
20 MLMAQEALA has no activity at all (Fig. 3b), suggesting that the C-terminal amino acids FL do significantly contribute to the recognition by the CTL.

Example 3

Comparison of CAMEL, LAGE-1^{S/L}, NY-ESO-1

As already mentioned, cDNA clone 4H8 lacks the first 84 bp of the LAGE-1^S
25 sequence, which means that it is devoid of the initiation codon at position 54 of that sequence (Fig. 2a). The first possible translation initiation site in 4H8 corresponds with the ATG at position 94 of LAGE-1^S, which is however, not in frame with the first ATG at position 54. Therefore, the protein translated from the 4H8 cDNA clone is different from the putative

LAGE-1 protein, since translation takes place in another reading frame (Fig. 2a and b). 4H8 encodes a protein of 109 amino acids (SEQ ID NO: 2) with a predicted molecular weight of 11.7 kD. The LAGE-1^S protein translated from the first ATG will be a 180 aa protein of 18.2 kD (SEQ ID NO: 4), while the unspliced variant, LAGE-1^L, encodes a 210 aa protein of 21.1 kD (SEQ ID NO: 6). NY-ESO-1 protein (SEQ ID NO: 8) is probably of the same size as LAGE-1^S, but differs at 26 amino acids. However, if translation of LAGE-1^{S/L} starts at the second ATG, proteins will be translated in another reading frame and are in that case identical to the protein translated from cDNA 4H8. Alternative translation of NY-ESO-1 (SEQ ID NO: 9 and NO: 10) results in a shorter variant of this protein (58 amino acids), because of an earlier stop codon (Fig. 2b), which differs from the CAMEL protein sequence only in its last 5 amino acids (Fig. 2b).

It was examined whether cells transfected with the complete LAGE-1 (or NY-ESO-1) cDNA clones are able to stimulate CTL 1/29. Remarkably, COS/HLA-A*0201 cells transfected with LAGE-1^S, the alternatively spliced LAGE-1^L (as well as with the NY-ESO-1) cDNA are able to stimulate CTL 1/29 (Fig. 4). This indicates that, at least in COS-7 cells, protein translation also starts from the second start codon at nucleotide 94 in LAGE-1^S, notwithstanding the presence of the first ATG at position 54. Also in this case, this results in the "alternative reading frame" peptide, MLMAQEALAF_L, recognized by CTL 1/29.

Example 4

Expression of CAMEL in E. Coli

To investigate whether CAMEL is indeed translated from the ORF-1 of the CAMEL (4H8) cDNA, the CAMEL cDNA (SEQ ID No: 1) was cloned in a bacterial expression vector (pET19b) (Studier et al., 1990). This vector contains a 6xHis-tag coding sequence, allowing detection of the His-tagged protein with an anti-His antibody. The pET19b-CAMEL construct was

transformed into E.coli and the bacteria were treated with IPTG to induce expression of the His-tagged CAMEL protein. Extracts were analyzed by Western blotting using the Penta-His antibody. Western blotting of a lysate shows a 15.5 kD protein, only slightly higher than the expected 14.5 kD of the His-tagged CAMEL protein after staining with a anti-His antibody (Fig. 5).

The CAMEL cDNA (SEQ ID No: 1) was cloned in pET19b and expressed in E.Coli. Lanes 1 and 2 represent the samples taken at 0h, lanes 3 and 4 at 4h after induction with IPTG. Because CAMEL might be an unstable protein, induction of protein expression was performed in the absence (lanes 1 and 3) or presence (lanes 2 and 4) of PMSF (a protease inhibitor). At the left the positions of the molecular weight marker proteins are indicated.

Example 5

Expression of LAGE-1 and NY-ESO-1 in healthy human tissues and melanoma cell lines

Hybridisation of Multiple Tissue Northern blots containing RNA of healthy human tissues with the LAGE-1^S cDNA showed high expression in testis and placenta and low, (but clear) expression in heart, skeletal muscle and pancreas (Fig. 6a). The positive signals exist of two bands, probably reflecting LAGE-1^S/NY-ESO-1 (750 bp) and LAGE-1^L (1000 bp).

Several melanoma cell lines were tested for expression of LAGE-1 and NY-ESO-1 by(Northern Blot analysis and) RT-PCR (Fig. 6b). Because of the strong homology between both genes, it is not possible to discriminate between LAGE-1 and NY-ESO-1 on Northern Blot. Therefore RT-PCR was performed with specific primers. In most cell lines a correlated expression of LAGE-1 and NY-ESO-1 was found; only cell line FM3.29 had expression of LAGE-1, but was negative for NY-ESO-1. Other cell lines expressed

either both or none of the two genes (Fig. 6b). There was a good correlation between the level of expression and the recognition by CTL 1/29 (Fig. 6b).

Example 6

5 Determination of LAGE1/CAMEL expression of in human tumors by RT-PCR

In order to evaluate the percentage of LAGE1/CAMEL positive human tumors, individual tumor tissues from breast or lung cancer patients were subjected to RT-PCR, as described in the Method section f), using the LAGE1-specific primers F3 (SEQ ID NO: 17) and ESO-1B (SEQ ID
10 NO: 18); reactions were performed in a Perkin Elmer 9600 thermocycler with 35 (instead of 30) cycles.

As shown in Table 1, approximately 50% (3/6) of the tested breast cancer specimens and 80% (10/12) of the lung cancer specimens were shown to be positive for LAGE1/CAMEL mRNA .

Table 1:

<i>Tumor type</i>	<i>#</i>	<i>LAGE1/CAMEL mRNA</i>
breast (ILC)	71526 / 85	-
breast (ILC)	44231 / 95	-
breast (ILC)	73507 / 95	+
breast (IDC)	19837 / 95	++
breast (IDC)	4385 / 95	+
breast (IDC)	48897 / 95	-
lung (AC)	15827 / 97	+
lung (AC)	53934 / 97	++
lung (AC)	67086 / 93	++
lung (AC)	62357 / 96	+
lung (AC)	T63244 / 93	-
lung (AC)	T64360 / 93	+
lung (SCC)	92710 / 96	++
lung (SCC)	53005	++
lung (SCC)	28649 / 97	-
lung (SCC)	16251 / 97	+
lung (SCC)	5063 / 93	++
lung (SCC)	7580 / 97	+
testis (positive control)	(GibcoBRL)	+

Example 7

Immunohistochemical analysis of CAMEL expression in human tumors

As described in Examples 5 and 6, LAGE-1 mRNA was detected in a panel
5 of tumor cell lines and tumor tissues and in a restricted number of healthy
tissues by means of RT-PCR and Northern blot experiments. However, it
remained to be determined whether in cells expressing the LAGE-1 mRNA
also the alternatively translated CAMEL is produced.

10 Frozen sections from a panel of different human tumors were analyzed by
immunohistochemistry using a CAMEL-specific rabbit antiserum B5 which
was affinity-purified against the B5 peptide. (For the preparation of the
antisera, see h) in the Methods section. The B5 antiserum was used
because B5 is CAMEL-specific, while the F4 antiserum may also recognize
an epitope present on a protein expressed from the ORF-1 of NY-ESO-1).
15 Specificity of the purified serum was demonstrated by peptide ELISA and
Western blotting against COS cells transfected with a CAMEL-GFP fusion
protein. Immunohistochemistry was performed using a 3-step avidin-biotin-
peroxidase staining procedure. The results are summarized in Table 2,
examples are shown in Figure 7.

Table 2:

Tumor type	Percentage of positive tumor cells				
(n=number of cases)	negative	<10%	10-40%	41-70%	>70%
Breast AC (n=11)	2	4	1	2	2
Colon AC (n=8)	1	0	1	1	5
Lung AC (n=10)	2	2	2	3	1
Lung SCC (n=9)	4	3	1	1	0
Pancreas AC (n=10)	1	2	0	2	5

AC: adenocarcinoma

SCC: squamous cell carcinoma

5

A total of 48 different specimen was investigated. In the majority of cases (38/48) CAMEL expression was detected. About half of the positive cases showed expression of CAMEL in 40% or more of the tumor cells, in some of the cases close to 100% of the tumor cells showed CAMEL-specific staining (an example is shown Figure 7, Colon AC). In the majority of tumor specimens expression was heterogeneous ranging from less than 10% of positive tumor cells to more than 70% of positive tumor cells (Table 2; examples are shown in Figure 7, arrows indicate positive tumor cell staining).

10

15 Example 8

Identification of HLA-A2 binding peptides within the CAMEL ORF

In order to identify further HLA-A2 epitopes besides the CTL-epitope (M) LMAQEALAF L (SEQ ID NO:11 and 12), CAMEL (SEQ ID NO:2) was examined according to the algorithms and motifs published by Parker et al.;

1994, and Rammensee et al., 1995. The result of this analysis indicated that several further peptides within the CAMEL protein have the potential to bind to HLA-A2, and three of these candidates -CAMEL10: FLMAQGAML (SEQ ID NO: 24), CAMEL16: AMLAAQERRV (SEQ ID NO: 25) and
5 CAMEL17: MLAAQERRV (SEQ ID NO:26) – were synthesized.

These synthetic peptides were evaluated for their ability to increase surface HLA-A2 expression on the transport defective cell line 174CEM.T2 (Nijman et al., 1993). Briefly, 5×10^5 cells/0.2 ml/well were seeded in 96-well V-bottom plates and incubated for 16 hours with increasing amounts
10 (0-320 µg/ml) of peptide at 37°C in a humidified atmosphere. HLA-A2 surface expression was measured by FACS analysis (Becton Dickinson) using purified BB7.2 as primary antibody and a goat-anti-mouse IgG RPE conjugate (DAKO) as detection antibody.

As positive controls the known HLA-A2 restricted CTL-epitopes from
15 CAMEL (MLMAQEALAF, SEQ ID NO:11) or tyrosinase (Wölfel et al., 1994; YMNGTMSQV, SEQ ID NO:27) were applied. Negative controls included an HLA-A1 binding (and therefore irrelevant) peptide from MAGE-3 (Gaugler et al. , 1994; EVDPIGHLY, SEQ ID NO:28) or no peptide at all.

20 The results from these experiments suggest that the nonamer CAMEL10 binds with similar affinity to HLA-A2 as compared to the positive controls used in the assay. The two other peptides (CAMEL16 and CAMEL17) showed only low affinity in this assay. Therefore in particular CAMEL10 represents a potential new HLA-A2 restricted CTL-epitope derived from
25 CAMEL protein (FIG. 8).

The testing of the immunogenicity of CAMEL10 and if it represents a naturally processed and presented ligand can be done as described in WO 97/30721 and Schweighoffer, 1997.

Bibliography

- Bakker, A.B.H., van der Burg, S.H., Huijbens, R.J.F., Drijfhout, J.-W.,
5 Melief, J.M., Adema, G.J., and Figdor, C.G. (1997). Analogues of CTL
epitopes with improved MHC Class-I Binding capacity elicit anti-melanome
CTL recognizing the wild-type epitope. *Int. J. Cancer* 70, 302-309.
- Blake, J., Johnston, J.V., Hellström, K.E., Marquardt, H., and, Chen, L.
10 (1996). Use of Combinatorial Peptide Libraries to Construct
Functional Mimics of Tumor Epitopes Recognized by MHC Class I-
Restricted Cytolytic T Lymphocytes. *J. Exp. Med.* 184, 121-130.
- Chen, Y.T., Scanlan, M.J., Sahin, U., Tureci, O., Gure, A.O., Tsang, S.,
15 Williamson, B., Stockert, E., Pfrenders, M., and Old, L.J. (1997). A
testicular antigen aberrantly expressed in human cancers detected by
autologous antibody screening. *Proc. Natl. Acad. Sci. U. S. A.* 94,
1914-1918.
- 20 Coulie P.G. (1997) Human tumour antigens recognized by T cells: new
perspectives for anti cancer vaccines. *Mol. Med. Today* 3, 261-268
- D'Amaro, J., Houbiers, J.G., Drijfhout, J.W., Brandt, R.M., Schipper, R.,
Bavinck, J.N., Melief, C.J., and Kast, W.M. (1995). A computer program for
25 predicting possible cytotoxic T lymphocyte epitopes based on HLA class I
peptide-binding motifs. *Hum. Immunol.* 43, 13-18.
- Dranoff, G., Jaffee, E., Lazenby, A., Golumbek, P., Levitsky, H., Brose, K.,
Jackson, V., Hamada, H., Pardoll, D., and Mulligan, R. (1993). Vaccination
30 with irradiated tumor cells engineered to secrete murine
granulocyte-macrophage colony-stimulating factor stimulates potent,

specific, and long-lasting anti-tumor immunity. *Proc. Natl. Acad. Sci. USA* **90**, 3539-3543.

Drijfhout, J.W., Brandt, R.M., D'Amaro, J., Kast, W.M., and Melief, C.J.
5 (1995). Detailed motifs for peptide binding to HLA-A*0201 derived from large random sets of peptides using a cellular binding assay. *Hum. Immunol.* **43**, 1-12.

Falk, K., Rötzschke, O., Stevanović, S., Jung, G., and Rammensee, H-G
10 (1991). Allele-specific motifs revealed by sequencing of self-peptides eluted from MHC molecules. *Nature* **351**, 290-296.

Fearon, E.R., Pardoll, D.M., Itaya, T., Golumbek, P., Levitsky, H.I., Simons, J.W., Karasuyama, H., Vogelstein, B., and Frost, P. (1990). Interleukine-2
15 production by tumor cells bypasses T helper function in the generation of an antitumor response. *Cell* **60**, 397-403.

Gansbacher, B., Zier, K., Daniels, B., Cronin, K., Bannerji, R., and Gilboa, E. (1990). Interleukin 2 gene transfer into tumor cells abrogates
20 tumorigenicity and induces protective immunity. *J. Exp. Med.* **172**, 1217-1224.

Gaugler B, van den Eynde B, van der Bruggen P, Romero P, Gaforio JJ, De Plaen E, Lethe B, Brasseur F, Boon T (1994) *J. Exp. Med.* **179**:921-30
25

Jäger, E., Chen, Y.T., Drijfhout, J.W., Karbach, J., Ringhoffer, M., Jäger, D., Arand, M., Wada, H., Noguchi, Y., Stockert, E., Old, L.J., and Knuth, A. (1998). Simultaneous humoral and cellular immune response against cancer-testis antigen NY-ESO-1: definition of human histocompatibility
30 leukocyte antigen (HLA)-A2-binding peptide epitopes. *J. Exp. Med.* **187**, 265-270.

- Kawakami, Y., Eliyahu, S., Jennings, C., Sakaguchi, K., King, X., Southwood, S., Robbins, P.F., Sette, A., Appella, E., and Rosenberg, S.A. (1995). Recognition of multiple epitopes in the human melanoma antigen gp100 by tumor-infiltrating T lymphocytes associated with *in vivo* tumor regression. *J. Immunol.* **154**, 3961-3968.
- 5 Lethe, B., Lucas, S., Michaux, L., Desmet, C., Godelaine, D., Serrano, A., Deplaen, E. and Boon, T. (1998). LAGE-1, a new gene with tumor specificity. *Int. J. Cancer* **76** (6):903-908.
- 10 Lin, A.Y., Devaux, B., Green, A., Sagerström, C., Elliott, J.F., and Davis, M.M. (1990). Expression of T Cell Antigen Receptor Heterodimers in a Lipid-Linked Form. *Science* **249**, 677-679.
- 15 Mayordomo, J.I., Zorina, T., Storkus, W.J., Zitvogel, L., Celluzzi, C., Falo, L.D., Melief, C.J., Ildstad, S.T., Kast, W.M., DeLeo, A.B., and Lotze, M.T. (1995). Bone marrow-derived dendritic cells pulsed with synthetic tumour peptides elicit protective and therapeutic antitumour immunity. *Nature Medicine* **1**, 1297-1302.
- 20 Nijman HW, Houbiers JG, Vierboom MP, van der Burg SH, Drijfhout JW, D'Amato J, Kenemans P, Melief CJ, Kast WM (1993) *Eur. J. Immunol.* **6**: 1215-9
- 25 Osanto, S., Brouwenstyn, N., Vaessen, N., Figdor, C.G., Melief, C.J., and Schrier, P.I. (1993). Immunization with interleukin-2 transfected melanoma cells. A phase I-II study in patients with metastatic melanoma. *Hum. Gene Ther.* **4**, 323-330.
- 30 Parker KC, Bednarek MA, Coligan JE (1994), *J. Immunol.* **152**:163

- Parkhurst, M.R., Salgaller, M.L., Southwood, S., Robbins, P.F., Sette, A., Rosenberg, S.A., and Kawakami, Y. (1996). Improved induction of melanoma-reactive CTL with peptides from the melanoma antigen gp100 modified at HLA-A*0201-binding residues. *J. Immunol.* **157**, 2539-2548.
- 5 Rammensee, H.G. and Friede, T. (1995). MHC ligands and peptide motifs: first listing. *Immunogenetics* **41**, 179-228.
- Ruppert, J., Sidney, J., Celis, E., Kubo, R.T., Grey, H.M., and Sette, A. (1993). Prominent role of secondary anchor residues in peptide binding to HLA-A2.1 molecules. *Cell* **74**, 929-937.
- 10 Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular Cloning, a Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press).
- 15 Schweighoffer, T. (1997) Molecular cancer vaccines: tumor therapy using antigen-specific immunizations. *Path. Onc. Res.* **3**, 164-176
- Sette, A., Vitiello, A., Reheman, B., Fowler, P., Nayersina, R., Kast, W.M., Melief, C.J.M., Oseroff, C., Yuan, L., Ruppert, J., Sidney, J., del Guercio, M.-F., Southwood, S., Kubo, R.T., Chesnut, R.W., Grey, H.M., and Chisari, F.V. (1994). The relationship between class I binding affinity and immunogenicity of potential cytotoxic T Cell epitopes. *J. Immunol.* **153**, 5586-5592.
- 20 Stauss, H.J., Davies, H., Sadovnikova, E., Chain, B., Horowitz., and Sinclair, C. (1992). Induction of cytotoxic T lymphocytes with peptides *in vitro*: Identification of candidate T-cell epitopes in human papilloma virus. *Proc. Natl. Acad. Sci. USA.* **89**, 7871-7875.
- 25

Studier, F.W., Rosenberg, A.H., Dunn, J.J. and Dubendorf, J.W. (1990).
Use of T7 RNA polymerase to direct expression of cloned genes. *Methods*
in *Enzymology* 185, 60-89

- 5 Tepper, R.I., Pattengale, P.K., and Leder, P. (1989). Murine interleukine-4
displays potent anti-tumour activity *in vivo*. *Cell* 57, 503-512.

Toes, R.E., Hoebe, R.C., Van der Voort, E., Rensing, M.E., Van-der-Eb,
A.J. Melief, C.J.M., and Offringa, R. 1997 Protective anti-tumor immunity
10 induced by vaccination with recombinant adenoviruses encoding multiple
tumor-associated cytotoxic T lymphocyte epitopes in a string-of-beads
fashion. *Proc. Natl. Acad. Sci. U.S.A.* 94 (26):14660-14665

Tuting, T., DeLeo, A.B., Lotze, M.T., and Storkus, W.J. Genetically modified
15 bone marrow-derived dendritic cells expressing tumor-associated viral or
"self" antigens induce antitumor immunity *in vivo*. *Eur. J. Immunol.* 27,
2702-2707, 1997.

van der Burg, S.H., et al., (1996), Immunogenicity of peptides bound to
20 MHC class I molecules depends on the MHC-peptide complex stability.
J. Immunol. 156, 3308-3314

Van den Eynde, B. and Brichard, V.G. (1995). New tumor antigens
recognized by T cells. *Current Opinion in Immunology* 7, 674-681.

25 van Elsas, A., van der Minne, C.E., Borghi, M., van der Spek, C.W.,
Braakman, E., Osanto, S., and Schrier, P.I. (1996). CTL Recognition of an
IL-2 Producing Melanoma Vaccine. In: *Immunology of human melanoma.*
Tumor-host interaction and immunotherapy, edited by M. Maio, Amsterdam:
30 IOS, 1996, p. 165-173.

- van Elsas, A., Aarnoudse, C., van der Minne, C.E., van der Spek, C.W.,
Brouwenstijn, N., Osanto, S., and Schrier, P.I. (1997). Transfection of IL-2
augments CTL response to human melanoma cells *in vitro*: immunological
characterization of a melanoma vaccine. *Journal of Immunotherapy* 20,
5 343-353.
- Versteeg, R., Noordermeer, I.A., Krüse-Wolters, K.M., Ruiter, D.J., and
Schrier, P.I. (1988). c-Myc downregulates class I HLA expression in human
melanomas. *EMBO J.* 7, 1023-1029.
- 10 Wang, R.F., Johnston, S.L., Zeng, G., Topalian, S.L., Schwartzentruber, D.J.,
and Rosenberg, S.A. (1998). A breast and melanoma-shared tumor antigen:
T cell responses to antigenic peptides translated from different open
reading frames. *J. Immunol.* 161, 3596-3606.
- 15 Wölfel T, van Pel A, Brichard V, Schneider J, Seliger B, Meyer zum
Büschhof KH, Boon T (1994) *Eur. J. Immunol.* 24:759-64
- 20 Wu, T.C., Guarnieri, F.G., Staveley-O'Carroll, K.F., Viscidi, R.P., Levitsky,
H.I., Hedrick, L., Cho, K.R., August, J.T., and Pardoll, D.M. (1995).
Engineering an intracellular pathway for major histocompatibility complex
class II presentation of antigens. *Proc. Natl. Acad. Sci. U.S.A.* 92(25):
11671-11675.
- 25 Zitvogel, L., Mayordomo, J.I., Tjandrawan, T., DeLeo, A.B., Clarke, M.R.,
Lotze, M.T., and Storkus, W.J. (1996). Therapy of murine tumors with tumor
peptide-pulsed dendritic cells: dependence on T cells, B7 costimulation,
and T helper cell 1-associated cytokines. *J. Exp. Med.* 183, 87-97.


Claims

1. Tumor-associated antigen CAMEL which is encoded by the ORF-1 of LAGE-1 cDNA and has the amino acid sequence set forth in
5 SEQ ID NO: 2.
2. The tumor-associated antigen of claim 1 for use in cancer therapy.
3. Immunogenic (poly)peptide derived from the tumor-associated antigen of claim 1.
4. An immunogenic peptide of claim 3, characterized in that it has the
10 amino acid sequence Met Leu Met Ala Gln Glu Ala Leu Ala Phe Leu (SEQ ID NO: 11).
5. An immunogenic peptide of claim 3, characterized in that it has the amino acid sequence Leu Met Ala Gln Glu Ala Leu Ala Phe Leu (SEQ ID NO: 12).
- 15 6. An immunogenic peptide of claim 3, characterized in that it has the amino acid sequence Phe Leu Met Ala Gln Gly Ala Met Leu (SEQ ID NO: 24).
7. An immunogenic peptide of claim 3, characterized in that it has the amino acid sequence Ala Met Leu Ala Ala Gln Glu Arg Arg Val
20 (SEQ ID NO: 25).
8. An immunogenic peptide of claim 3, characterized in that it has the amino acid sequence Met Leu Ala Ala Gln Glu Arg Arg Val (SEQ ID NO: 26).
9. An immunogenic peptide of any one of claims 3 to 8 for use in cancer
25 immunotherapy.
10. Pharmaceutical composition containing the tumor-associated antigen CAMEL of claim 1.

11. Pharmaceutical composition containing an immunogenic peptide of any one of claims 3 to 8.
12. Isolated DNA molecule comprising the sequence set forth in SEQ ID NO: 1.
- 5 13. Recombinant DNA molecule comprising the DNA molecule of claim 12.
14. A DNA molecule of claim 12 or 13 for use in cancer immunotherapy.

1/15

Fig. 1



2/15
Fig. 2A

FIGURE 2 A)

CAMEL	-----	
LAGE-1 ^S	--ATCCTCGTGGCCCTGACCTTCTCTGAGAGCCGGGCAGAGGCTCCG	48
LAGE-1 ^L	GATCCTCGTGGCCCTGACCTTCTCTGAGAGCCGGGCAGAGGCTCCG	50
NY-ESO-1	--ATCCTCGTGGCCCTGACCTTCTCTGAGAGCCGGGCAGAGGCTCCG	48
CAMEL	-----	
LAGE-1 ^S	GAGCC ATG CAGGCCGAAGGCCAGGCCACAGGGGTTTCGACGGCGATGCT	14
LAGE-1 ^L	GAGCC ATG CAGGCCGAAGGCCAGGCCACAGGGGTTTCGACGGCGATGCT	98
NY-ESO-1	GAGCC ATG CAGGCCGAAGGCCAGGGGTTTCGACGGCGATGCT	100
	*****	98
CAMEL		
LAGE-1 ^S	GATGGCCAGGAGGCCCTGGCATTCCTGATGGCCAGGGGGCAATGCTGG	64
LAGE-1 ^L	GATGGCCAGGAGGCCCTGGCATTCCTGATGGCCAGGGGGCAATGCTGG	148
NY-ESO-1	GATGGCCAGGAGGCCCTGGCATTCCTGATGGCCAGGGGGCAATGCTGG	150
	*****	148
CAMEL		
LAGE-1 ^S	CGGCCAGGAGAGGCGGGTGCCACGGGGCGGCAGAGTCCCCGGGGCGCAG	114
LAGE-1 ^L	CGGCCAGGAGAGGCGGGTGCCACGGGGCGGCAGAGTCCCCGGGGCGCAG	198
LAGE-1 ^L	CGGCCAGGAGAGGCGGGTGCCACGGGGCGGCAGAGTCCCCGGGGCGCAG	200
LAGE-1 ^L	CGGCCAGGAGAGGCGGGTGCCACGGGGCGGCAGAGTCCCCGGGGCGCAG	198

Fig. 2A continued

CAMEL
 LAGE-1^S
 LAGE-1^L
 NY-ESO-1

GGCAGCAAGGCCCTCGGGCCGAGAGGCGCCCGGGGTCCGCAT
 GGCAGCAAGGCCCTCGGGCCGAGAGGCGCCCGGGGTCCGCAT
 GGCAGCAAGGCCCTCGGGCCGAGAGGCGCCCGGGGTCCGCAT
 GGCAGCAAGGCCCTCGGGCCGAGAGGCGCCCGGGGTCCGCAT

164
 248
 250
 248

CAMEL
 LAGE-1^S
 LAGE-1^L
 NY-ESO-1

GGCGTGCCGCTTCTGCGCAGGATGGAAGTGCCCTCGGGGCCAGGAG
 GGCGTGCCGCTTCTGCGCAGGATGGAAGTGCCCTCGGGGCCAGGAG
 GGCGTGCCGCTTCTGCGCAGGATGGAAGTGCCCTCGGGGCCAGGAG
 GGCGTGCCGCTTCTGCGCAGGATGGAAGTGCCCTCGGGGCCAGGAG

214
 298
 300
 298

CAMEL
 LAGE-1^S
 LAGE-1^L
 NY-ESO-1

GCCGGACAGCCGCTGCTTCAGTTGCACATCACGATGCCCTTCTCGTCGC
 GCCGGACAGCCGCTGCTTCAGTTGCACATCACGATGCCCTTCTCGTCGC
 GCCGGACAGCCGCTGCTTCAGTTGCACATCACGATGCCCTTCTCGTCGC
 GCCGGACAGCCGCTGCTTCAGTTGCACATCACGATGCCCTTCTCGTCGC

264
 348
 350
 348

CAMEL
 LAGE-1^S
 LAGE-1^L
 NY-ESO-1

CCATGGAAGCGGAGCTGGTCCGCAGGATCCTGTCCCGGGATGCCGCACCT
 CCATGGAAGCGGAGCTGGTCCGCAGGATCCTGTCCCGGGATGCCGCACCT
 CCATGGAAGCGGAGCTGGTCCGCAGGATCCTGTCCCGGGATGCCGCACCT
 CCATGGAAGCAGAGCTGGCCCGCAGGAGCCTGGCCAGGATGCCCCACCG

314
 398
 400
 398

Fig. 2A continued

CAMEL	CTCCCCGACCAAGGGCGGTTCTGAAGGACTTCACCGTGTCCGGCAACCT	364
LAGE-1 ^s	CTCCCCGACCAAGGGCGGTTCTGAAGGACTTCACCGTGTCCGGCAACCT	448
LAGE-1 ^L	CTCCCCGACCAAGGGCGGTTCTGAAGGACTTCACCGTGTCCGGCAACCT	450
NY-ESO-1	CTTCCCGTGCCAGGGGTGCTTCTGAAGGAGTTCACTGTGTCCGGCAACAT	448
	** *** . . ***** * ***** * ***** *	
CAMEL	ACTGTTTAT-----	373
LAGE-1 ^s	ACTGTTTAT-----	457
LAGE-1 ^L	ACTGTTTATGTCAGTTCGGGACCAGGACAGGGAAGCGCTGGGCGGATGA	500
NY-ESO-1	ACTGACTAT-----	457
	****. ***	
CAMEL	-----	373
LAGE-1 ^s	-----	457
LAGE-1 ^L	GGTGTTGGGCTGGGATCCGCCCTCCCCGAGGGGCAGAAAGCT	550
NY-ESO-1	-----	457
CAMEL	-----	373
LAGE-1 ^s	-----	457
LAGE-1 ^L	AGAGATCTCAGAACACCCCAACACAAGGTCTCAGAACAGAGACCTGGTAC	600
NY-ESO-1	-----	457

Fig. 2A continued

```
373  -----
457  -----
650  ACCAGGCCCGCCGCCACCCGAGGAGCCAGGAGATGGGTGCAGAGGTG
457  -----

385  -----CCGACTGACTGC
469  -----CCGACTGACTGC
700  TCGCCTTTAATGTGATGTTCTCTGCCCCCTCACATTTAGCCGACTGACTGC
469  -----CCGACTGACTGC
      *****

435  TGCAGACCACCGCCAACTGCAGCTCTCCATCAGCTCCTGTCTCCAGCAGC
519  TGCAGACCACCGCCAACTGCAGCTCTCCATCAGCTCCTGTCTCCAGCAGC
750  TGCAGACCACCGCCAACTGCAGCTCTCCATCAGCTCCTGTCTCCAGCAGC
519  TGCAGACCACCGCCAACTGCAGCTCTCCATCAGCTCCTGTCTCCAGCAGC
      *****

485  TTTCCCTGTTGATGTGGATCACGCAGTGCTTTCTGCCCGTGTTTTGGCT
569  TTTCCCTGTTGATGTGGATCACGCAGTGCTTTCTGCCCGTGTTTTGGCT
800  TTTCCCTGTTGATGTGGATCACGCAGTGCTTTCTGCCCGTGTTTTGGCT
569  TTTCCCTGTTGATGTGGATCACGCAGTGCTTTCTGCCCGTGTTTTGGCT
      *****
```

CAMEL
LAGE-1^s
LAGE-1^L
NY-ESO-1

CAMEL
LAGE-1^s
LAGE-1^L
NY-ESO-1

CAMEL
LAGE-1^s
LAGE-1^L
NY-ESO-1

CAMEL
LAGE-1^s
LAGE-1^L
NY-ESO-1

6/15

Fig. 2A continued

CAMEL
 LAGE-1^s
 LAGE-1^L
 NY-ESO-1

CAGGCTCCCTCAGGGCAGAGCGCTAAGCCAGCCTGGCGCCCTTCCTA 535
 CAGGCTCCCTCAGGGCAGAGCGCTAAGCCAGCCTGGCGCCCTTCCTA 619
 CAGGCTCCCTCAGGGCAGAGCGCTAAGCCAGCCTGGCGCCCTTCCTA..850
 CAGCCTCCCTCAGGGCAGAGCGCTAAGCCAGCCTGGCGCCCTTCCTA..519

CAMEL
 LAGE-1^s
 LAGE-1^L
 NY-ESO-1

GGTCATGCCCTCCTCCCTAGGGAATGGTCCAGCACGAGTGGCCAGTTCA 585
 GGTCATGCCCTCCTCCCTAGGGAATGGTCCAGCACGAGTGGCCAGTTCA 669
 GGTCATGCCCTCCTCCCTAGGGAATGGTCCAGCACGAGTGGCCAGTTCA 900
 GGTCATGCCCTCCTCCCTAGGGAATGGTCCAGCACGAGTGGCCAGTTCA 669

CAMEL
 LAGE-1^s
 LAGE-1^L
 NY-ESO-1

TTGTGGGGCCCTGATTGTTGTCTCGCTGGAGGAGGCGCTTACATGTTG..635
 TTGTGGGGCCCTGATTGTTGTCTCGCTGGAGGAGGCGCTTACATGTTG..719
 TTGTGGGGCCCTGATTGTTGTCTCGCTGGAGGAGGCGCTTACATGTTG..950
 TTGTGGGGCCCTGATTGTTGTCTCGCTGGAGGAGGCGCTTACATGTTG..719

CAMEL
 LAGE-1^s
 LAGE-1^L
 NY-ESO-1

TTTCTGTAGAAAATAAAGCTGAGCTACGAAAAAATAAAAAA-----679
 TTTCTGTAGAAAATAAAGCTGAGCTACGAAAAAATAAAAAA-----767
 TTTCTGTAGAAAATAAAGCTGAGCTACGAAAAAATAAAAAA-----993
 TTTCTGTAGAAAATAAAGCTGAGCTACGAAAAAATAAAAAA-----752

7/15

Fig. 2B

Protein Translations

ORF3

LAGE-1^S MQAEGQGTGGSTGDADGPGGGPGIPDGPGGNAGGPGEAGAT 40
 LAGE-1^L MQAEGQGTGGSTGDADGPGGGPGIPDGPGGNAGGPGEAGAT 40
 NY-ESO-1 MQAEGRGTGGSTGDADGPGGGPGIPDGPGGNAGGPGEAGAT 40

LAGE-1^S GGRGPRGAGAAARASGPRGGAPRGPHGGAASAQDGRCPCCA 80
 LAGE-1^L GGRGPRGAGAAARASGPRGGAPRGPHGGAASAQDGRCPCCA 80
 NY-ESO-1 GGRGPRGAGAAARASGPGGGAPRGPHGGAASGLNGCCRCGA 80

LAGE-1^S RRPDSRLLQLHITMPFSSPMEAEELVRRILSRDAAPLPRPG 120
 LAGE-1^L RRPDSRLLQLHITMPFSSPMEAEELVRRILSRDAAPLPRPG 120
 NY-ESO-1 RGPEsrLL**EFY**LAMP**FA**TPMEAEELARRSL**AQ**DAPPL**VP**PG 120

LAGE-1^S AVLKDFTVSGNLLFIRLTAADHRQLQLSISSCLQQLSLLM 160
 LAGE-1^L AVLKDFTVSGNLL**FMSVRDQDREGAGRMRVV**GWGLGS**ASP** 160
 NY-ESO-1 **VLLK**EFTVSGN**IL**TIRLTAADHRQLQLSISSCLQQLSLLM 160


8/15

Fig. 2B continued

LAGE-1 ^S	WITQCFLPVFLAQAPSGQRR	180
LAGE-1 ^L	EGQKARDLRTPKHKVSEQRPGTGPFPPEGAQGDGCRGVA	200
NY-ESO-1	WITQCFLPVFLAQPPSGQRR	180
LAGE-1 ^S		180 aa, 18.2 kD
LAGE-1 ^L	FNVMFSAPHI	210 aa, 21.1 kD
NY-ESO-1		180 aa, 18.2 kD
ORF1		
LAGE-1 ^S	<u>MLMAQEALAF</u> MLMAQGAMLAQAQERRVPRAAEVPGAQGGQGP	40
LAGE-1 ^L	<u>MLMAQEALAF</u> MLMAQGAMLAQAQERRVPRAAEVPGAQGGQGP	40
NY-ESO-1	<u>MLMAQEALAF</u> MLMAQGAMLAQAQERRVPRAAEVPGAQGGQGP	40
LAGE-1 ^S	RGREEAPRGVRMAVPLLRMEGAPAGPGGRTAACFSCTSR	80
LAGE-1 ^L	RGREEAPRGVRMAVPLLRMEGAPAGPGGRTAACFSCTSR	80
NY-ESO-1	RGREEAPRGVRMA ARLQG	58
LAGE-1 ^S	CLSRRPWKRWSAGSCPGMPHLSPDQGRF	109 aa, 11.7 kD
LAGE-1 ^L	CLSRRPWKRWSAGSCPGMPHLSPDQGRF	109 aa, 11.7 kD
NY-ESO-1		58 aa, 6.2 kD


9/15

Fig. 3




10/15

Fig. 4




11/15

Fig. 5



12/15

Fig. 6A



13/15

Fig. 6B






Fig. 7



15/15

Fig. 8



SEQUENCE LISTING

5

(1) GENERAL INFORMATION:

10

(i) APPLICANT:

- (A) NAME: Boehringer Ingelheim International GmbH
- (B) STREET: Binger Strasse 173
- (C) CITY: Ingelheim am Rhein
- (E) COUNTRY: Germany
- (F) POSTAL CODE (ZIP): 55216
- (G) TELEPHONE: 06132/772282
- (H) TELEFAX: 06132/774377

15

20

(ii) TITLE OF INVENTION: Tumor-associated Antigen

(iii) NUMBER OF SEQUENCES: 28

25

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

30

35

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 679 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: cDNA to mRNA

45

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

50

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: homo sapiens
- (F) TISSUE TYPE: Melanoma

(ix) FEATURE:

- (A) NAME/KEY: 3'UTR

55

(B) LOCATION:340..679

(ix) FEATURE:

5 (A) NAME/KEY: 5'UTR
(B) LOCATION:1..9

(ix) FEATURE:

10 (A) NAME/KEY: CDS
(B) LOCATION:10..339

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

15	CGACGGGCG ATG CTG ATG GGC CAG GAG GGC CTG GCA TTC CTG ATG GGC Met Leu Met Ala Gln Glu Ala Leu Ala Phe Leu Met Ala 1 5 10	48
20	CAG GGG GCA ATG CTG GGG GGC CAG GAG AGG GGG GTG CCA GGG GGG GCA Gln Gly Ala Met Leu Ala Ala Gln Glu Arg Arg Val Pro Arg Ala Ala 15 20 25	96
25	GAG GTC CCC GGG GGG CAG GGG CAG CAA GGG OCT GGG GGC CGA GAG GAG Glu Val Pro Gly Ala Gln Gly Gln Gln Gly Pro Arg Gly Arg Glu Glu 30 35 40 45	144
30	GGG CCC GGC GGG GTC GGC ATG GGG GTG CCG CTT CTG CCC AGG ATG GAA Ala Pro Arg Gly Val Arg Met Ala Val Pro Leu Leu Arg Arg Met Glu 50 55 60	192
35	GGT GGC OCT GGG GGG CCA GGA GGC GGG ACA GGC GGC TGC TTC AGT TGC Gly Ala Pro Ala Gly Pro Gly Gly Arg Thr Ala Ala Cys Phe Ser Cys 65 70 75	240
40	ACA TCA CGA TGC CTT TCT GGT GGC CCA TGG AAG GGG AGC TGG TCC GCA Thr Ser Arg Cys Leu Ser Arg Arg Pro Trp Lys Arg Ser Trp Ser Ala 80 85 90	288
45	GGA TCC TGT CCC GGG ATG CCG CAC CTC TCC CCC GAC CAG GGG GGG TTC Gly Ser Cys Pro Gly Met Pro His Leu Ser Pro Asp Gln Gly Arg Phe 95 100 105	336
50	TGA AGGACTTCAC CGTGTCCGGC AACCTACTGT TTATCCGACT GACTGCTGCA *	389
55	110	
	GACCAACGGC AACTGCAGCT CTCACACAGC TCTGTCTCC AGCAGCTTTC CCTGTGTATG	449
	TGGATCAGGC AGTGCCTTCT GCGGTGTATT TTGGCTCAGG CTCCTCAGG GCAGAGGGCC	509
	TAAGGACAGC CTGGGCGGCC TTCTAGGTC ATGCTCTCTC CCTAGGGAA TGGTCCAGC	569
	ACGAGTGGCC AGTTCATTGT GGGGGCTGA TTGTTGTGCG CTGAGGAGG ACGGCTTACA	629
	TGTTTGTTC TGTAGAAAT AAAGCTGAGC TACGAAAAA AAAAAAAAAA	679

(2) INFORMATION FOR SEQ ID NO: 2:

5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 109 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

15 Met Leu Met Ala Gln Glu Ala Leu Ala Phe Leu Met Ala Gln Gly Ala
 1 5 10 15

Met Leu Ala Ala Gln Glu Arg Arg Val Pro Arg Ala Ala Glu Val Pro
 20 25 30

20 Gly Ala Gln Gly Gln Gln Gly Pro Arg Gly Arg Glu Glu Ala Pro Arg
 35 40 45

Gly Val Arg Met Ala Val Pro Leu Leu Arg Arg Met Glu Gly Ala Pro
 50 55 60

25 Ala Gly Pro Gly Gly Arg Thr Ala Ala Cys Phe Ser Cys Thr Ser Arg
 65 70 75 80

30 Cys Leu Ser Arg Arg Pro Trp Lys Arg Ser Trp Ser Ala Gly Ser Cys
 85 90 95

Pro Gly Met Pro His Leu Ser Pro Asp Gln Gly Arg Phe *
 100 105 110

35

(2) INFORMATION FOR SEQ ID NO: 3:

40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 767 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

50 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: homo sapiens
 (F) TISSUE TYPE: Melanoma

55 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 54..596

(ix) FEATURE:

(A) NAME/KEY: 3'UTR

(B) LOCATION:597..767

5

(ix) FEATURE:

(A) NAME/KEY: 5'UTR

(B) LOCATION:1..53

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

	ATCTCTGGTGG GGGCTGAOCT TCTCTCTGAG AGGCGGCGAG AGGCTCGGA GCC ATG	56
		Met
15		1
	CAG GCC GAA GGC CAG GGC ACA GGG GGT TCG ACG GGC GAT GCT GAT GGC	104
	Gln Ala Glu Gly Gln Gly Thr Gly Gly Ser Thr Gly Asp Ala Asp Gly	
	5 10 15	
20	CCA GGA GGC OCT GGC ATT OCT GAT GGC CCA GGG GGC AAT GCT GGC GGC	152
	Pro Gly Gly Pro Gly Ile Pro Asp Gly Pro Gly Gly Asn Ala Gly Gly	
	20 25 30	
25	CCA GGA GAG GGG GGT GGC ACG GGC GGC AGA GGT CCC CCG GGC CCA GGC	200
	Pro Gly Glu Ala Gly Ala Thr Gly Gly Arg Gly Pro Arg Gly Ala Gly	
	35 40 45	
30	GCA CCA AGG GGC TCG GGG CCG AGA GGA GGC GGC CCG CCG GGT CCG CAT	248
	Ala Ala Arg Ala Ser Gly Pro Arg Gly Gly Ala Pro Arg Gly Pro His	
	50 55 60 65	
	GGC GGT GGC GCT TCT GCG CAG GAT GGA AGG TGC CCC TGC GGG GGC AGG	296
	Gly Gly Ala Ala Ser Ala Gln Asp Gly Arg Cys Pro Cys Gly Ala Arg	
35		70 75 80
	AGG CCG GAC AGC CCG CTG CTT CAG TTG CAC ATC ACG ATG OCT TTC TCG	344
	Arg Pro Asp Ser Arg Leu Leu Gln Leu His Ile Thr Met Pro Phe Ser	
	85 90 95	
40	TCG CCC ATG GAA GCG GAG CTG GTC CCG AGG ATC CTG TCC CCG GAT GGC	392
	Ser Pro Met Glu Ala Glu Leu Val Arg Arg Ile Leu Ser Arg Asp Ala	
	100 105 110	
45	GCA OCT CTC CCC CGA CCA GGG GGG GTT CTG AAG GAC TTC ACC GTC TCC	440
	Ala Pro Leu Pro Arg Pro Gly Ala Val Leu Lys Asp Phe Thr Val Ser	
	115 120 125	
	GGC AAC CTA CTG TTT ATC CGA CTG ACT GCT GCA GAC CAC CCG CAA CTG	488
50	Gly Asn Leu Leu Phe Ile Arg Leu Thr Ala Ala Asp His Arg Gln Leu	
	130 135 140 145	
	CAG CTC TCC ATC AGC TCC TGT CTC CAG CAG CTT TCC CTG TTG ATG TGG	536
	Gln Leu Ser Ile Ser Ser Cys Leu Gln Gln Leu Ser Leu Leu Met Trp	
55		150 155 160

ATC AGC CAG TGC TTT CTG CCC GTG TTT TTG GCT CAG GCT CCC TCA GGG 584
 Ile Thr Gln Cys Phe Leu Pro Val Phe Leu Ala Gln Ala Pro Ser Gly
 165 170 175
 5 CAG AGC CGC TAA GGCAGGCTG GGGCCCTTC CTAGGTCATG OCTOCTOCCC 636
 Gln Arg Arg *
 180
 TAGGGAATGS TOCCAGCAAG AGTGGCCAGT TCATTGTGGG GGCTGATTG TTGTGGCTG 696
 10 GAGGAGGAGG GCTTACAATG TTGTTCTGT AGAAAATAAA GCTCAGCTAC GAAAAAATAA 756
 AAAAAAATAA A 767
 15

(2) INFORMATION FOR SEQ ID NO: 4:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 180 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 25 (ii) MOLECULE TYPE: protein
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

30 Met Gln Ala Glu Gly Gln Gly Thr Gly Gly Ser Thr Gly Asp Ala Asp
 1 5 10 15
 Gly Pro Gly Gly Pro Gly Ile Pro Asp Gly Pro Gly Gly Asn Ala Gly
 20 25 30
 35 Gly Pro Gly Glu Ala Gly Ala Thr Gly Gly Arg Gly Pro Arg Gly Ala
 35 40 45
 Gly Ala Ala Arg Ala Ser Gly Pro Arg Gly Gly Ala Pro Arg Gly Pro
 50 55 60
 40 His Gly Gly Ala Ala Ser Ala Gln Asp Gly Arg Cys Pro Cys Gly Ala
 65 70 75 80
 Arg Arg Pro Asp Ser Arg Leu Leu Gln Leu His Ile Thr Met Pro Phe
 45 85 90 95
 Ser Ser Pro Met Glu Ala Glu Leu Val Arg Arg Ile Leu Ser Arg Asp
 100 105 110
 50 Ala Ala Pro Leu Pro Arg Pro Gly Ala Val Leu Lys Asp Phe Thr Val
 115 120 125
 Ser Gly Asn Leu Leu Phe Ile Arg Leu Thr Ala Ala Asp His Arg Gln
 130 135 140
 55 Leu Gln Leu Ser Ile Ser Ser Cys Leu Gln Gln Leu Ser Leu Leu Met
 145 150 155 160

Trp Ile Thr Gln Cys Phe Leu Pro Val Phe Leu Ala Gln Ala Pro Ser
 165 170 175

5 Gly Gln Arg Arg *
 180

(2) INFORMATION FOR SEQ ID NO: 5:

10

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 993 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

20

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

25

- (A) ORGANISM: homo sapiens
 (F) TISSUE TYPE: Melanoma

(ix) FEATURE:

30

- (A) NAME/KEY: 5'UTR
 (B) LOCATION:1..55

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION:56..688

35

(ix) FEATURE:

- (A) NAME/KEY: 3'UTR
 (B) LOCATION:689..993

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

	GCATCCTCGT GGGGCTGAC CTCCTCTCTG AGAGGCTGGC AGAGGCTCG GAGCC ATG	58
		Met
		1
45	CAG GGC GAA GGC CAG GGC ACA GGG GGT TCG ACG GGC GAT GCT GAT GGC	106
	Gln Ala Glu Gly Gln Gly Thr Gly Gly Ser Thr Gly Asp Ala Asp Gly	
	5 10 15	
50	CCA GGA GGC OCT GGC ATT OCT GAT GGC CCA GGG GGC AAT GCT GGC GGC	154
	Pro Gly Gly Pro Gly Ile Pro Asp Gly Pro Gly Gly Asn Ala Gly Gly	
	20 25 30	
55	CCA GGA GAG GGG GGT GGC ACG GGC GGC AGA GGT CCG CCG GGC GCA GGC	202
	Pro Gly Glu Ala Gly Ala Thr Gly Gly Arg Gly Pro Arg Gly Ala Gly	
	35 40 45	
	GCA GCA AGG GGC TCG GGG CCG AGA GGA GGC GGC CCG CCG GGT CCG CAT	250

7/18

	Ala Ala Arg Ala Ser Gly Pro Arg Gly Gly Ala Pro Arg Gly Pro His	
	50 55 60 65	
5	GGC GGT GCC GCT TCT GCG CAG GAT GGA AGG TGC CCC TGC GGG GCC AGG Gly Gly Ala Ala Ser Ala Gln Asp Gly Arg Cys Pro Cys Gly Ala Arg	298
	70 75 80	
10	AGG CCG GAC AGC CGC CTG CTT CAG TTG CAC ATC ACG ATG OCT TTC TCG Arg Pro Asp Ser Arg Leu Leu Gln Leu His Ile Thr Met Pro Phe Ser	346
	85 90 95	
15	TOG CCC ATG GAA GCG GAG CTG GTC CGC AGG ATC CTG TCC CGG GAT GCC Ser Pro Met Glu Ala Glu Leu Val Arg Arg Ile Leu Ser Arg Asp Ala	394
	100 105 110	
	GCA CCT CTC CCC CGA CCA GGG GCG GTT CTG AAG GAC TTC ACC GTG TCC Ala Pro Leu Pro Arg Pro Gly Ala Val Leu Lys Asp Phe Thr Val Ser	442
	115 120 125	
20	GGC AAC CTA CTG TTT ATG TCA GTT CCG GAC CAG GAC AGG GAA GGC GCT Gly Asn Leu Leu Phe Met Ser Val Arg Asp Gln Asp Arg Glu Gly Ala	490
	130 135 140 145	
25	GGG CCG ATG AGG GTG GTG GGT TGG GGG CTG GGA TCC GCC TCC CCG GAG Gly Arg Met Arg Val Val Gly Trp Gly Leu Gly Ser Ala Ser Pro Glu	538*
	150 155 160	
30	GGG CAG AAA GCT AGA GAT CTC AGA ACA CCC AAA CAC AAG GTC TCA GAA Gly Gln Lys Ala Arg Asp Leu Arg Thr Pro Lys His Lys Val Ser Glu	586
	165 170 175	
35	CAG AGA CCT GGT ACA CCA GGC CCG CCG CCA CCC GAG GGA GCC CAG GGA Gln Arg Pro Gly Thr Pro Gly Pro Pro Pro Glu Gly Ala Gln Gly	634
	180 185 190	
	GAT GGG TGC AGA GGT GTC GCC TTT AAT GTG ATG TTC TCT GGC OCT CAC Asp Gly Cys Arg Gly Val Ala Phe Asn Val Met Phe Ser Ala Pro His	682
	195 200 205	
40	ATT TAG CCGACTGACT GCCTGAGACC ACCGCCAACT GCAGCTCTCC ATCAGCTCCT Ile *	738
	210	
45	GTCTCAGCA GCTTTCCCTG TTGATGIGGA TCAAGCAGTG CTTTCTGCC GGTGTTTGG	798
	CTCAGGCTCC CTCAGGCGAG AGGCGCTAAG CCGAGCCTGG CGGCCCTCC TAGGTCATGC	858
	CTCTCCCTT AGGGAATGCT CCGAGCAAGA GGGCCAGTT CATGTGGGG GCGTGATTGT	918
50	TTGTCCCTGG AGGAGGAAGG CTTACATGTT TGTTCCTGTA GAAATATAAG CTGAGCTACG	978
	AAAAAAAAAAAA	993
55		

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 210 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

10 Met Gln Ala Glu Gly Gln Gly Thr Gly Gly Ser Thr Gly Asp Ala Asp
 1 5 10 15

Gly Pro Gly Gly Pro Gly Ile Pro Asp Gly Pro Gly Gly Asn Ala Gly
 20 25 30

15 Gly Pro Gly Glu Ala Gly Ala Thr Gly Gly Arg Gly Pro Arg Gly Ala
 35 40 45

Gly Ala Ala Arg Ala Ser Gly Pro Arg Gly Gly Ala Pro Arg Gly Pro
 20 50 55 60

His Gly Gly Ala Ala Ser Ala Gln Asp Gly Arg Cys Pro Cys Gly Ala
 65 70 75 80

25 Arg Arg Pro Asp Ser Arg Leu Leu Gln Leu His Ile Thr Met Pro Phe
 85 90 95

Ser Ser Pro Met Glu Ala Glu Leu Val Arg Arg Ile Leu Ser Arg Asp
 100 105 110

30 Ala Ala Pro Leu Pro Arg Pro Gly Ala Val Leu Lys Asp Phe Thr Val
 115 120 125

Ser Gly Asn Leu Leu Phe Met Ser Val Arg Asp Gln Asp Arg Glu Gly
 35 130 135 140

Ala Gly Arg Met Arg Val Val Gly Trp Gly Leu Gly Ser Ala Ser Pro
 145 150 155 160

40 Glu Gly Gln Lys Ala Arg Asp Leu Arg Thr Pro Lys His Lys Val Ser
 165 170 175

Glu Gln Arg Pro Gly Thr Pro Gly Pro Pro Pro Glu Gly Ala Gln
 180 185 190

45 Gly Asp Gly Cys Arg Gly Val Ala Phe Asn Val Met Phe Ser Ala Pro
 195 200 205

His Ile *
 50 210

55

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 752 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

10 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Homo sapiens

15 (ix) FEATURE:
 (A) NAME/KEY: 5'UTR
 (B) LOCATION:1..53

20 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION:54..596

25 (ix) FEATURE:
 (A) NAME/KEY: 3'UTR
 (B) LOCATION:597..752

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

30	ATCTCTGTTGG GCGCTGACCT TCTCTCTGAG AGCGGGGCGAG AGGCTCCGGA GGC ATG	56
	Met	
	1	
35	CAG GGC GAA GGC CGG GGC ACA GGG GGT TCG ACG GGC GAT GCT GAT GGC	104
	Gln Ala Glu Gly Arg Gly Thr Gly Gly Ser Thr Gly Asp Ala Asp Gly	
	5 10 15	
40	CCA GGA GGC CCT GGC ATT CCT GAT GGC CCA GGG GGC AAT GCT GGC GGC	152
	Pro Gly Gly Pro Gly Ile Pro Asp Gly Pro Gly Gly Asn Ala Gly Gly	
	20 25 30	
45	CCA GGA GAG GCG GGT GGC ACG GGC GGC AGA GGT CCC CGG GGC CCA GGG	200
	Pro Gly Glu Ala Gly Ala Thr Gly Arg Gly Pro Arg Gly Ala Gly	
	35 40 45	
50	GCA GCA AGG GGC TCG GGG CCG GGA GGA GGC GGC CCG CGG GGT CCG CAT	248
	Ala Ala Arg Ala Ser Gly Pro Gly Gly Gly Ala Pro Arg Gly Pro His	
	50 55 60 65	
55	GGC GGC GCG GCT TCA GGG CTG AAT GGA TGC TGC AGA TGC GGC GGC AGG	296
	Gly Gly Ala Ala Ser Gly Leu Asn Gly Cys Cys Arg Cys Gly Ala Arg	
	70 75 80	
55	GGG CCG GAG AGC CGC CTG CTT GAG TTC TAC CTC GGC ATG CCT TTC GCG	344
	Gly Pro Glu Ser Arg Leu Leu Glu Phe Tyr Leu Ala Met Pro Phe Ala	
	85 90 95	
	ACA CCC ATG GAA GCA GAG CTG GGC CGC AGG AGC CTG GGC CAG GAT GGC	392

Thr Pro Met Glu Ala Glu Leu Ala Arg Arg Ser Leu Ala Gln Asp Ala
 100 105 110

5 OCA CCG CTT CCC GTG CCA GGG GTG CTT CTG AAG GAG TTC ACT GTG TOC 440
 Pro Pro Leu Pro Val Pro Gly Val Leu Leu Lys Glu Phe Thr Val Ser
 115 120 125

10 GGC AAC ATA CTG ACT ATC CGA CTG ACT GCT GCA GAC CAC CGC CAA CTG 488
 Gly Asn Ile Leu Thr Ile Arg Leu Thr Ala Ala Asp His Arg Gln Leu
 130 135 140 145

CAG CTC TCC ATC AGC TCC TGT CTC CAG CAG CTT TCC CTG TTG ATG TGG 536
 Gln Leu Ser Ile Ser Ser Cys Leu Gln Gln Leu Ser Leu Leu Met Trp
 150 155 160

15 ATC ACG CAG TGC TTT CTG CCC GTG TTT TTG GCT CAG CCT CCC TCA GGG 584
 Ile Thr Gln Cys Phe Leu Pro Val Phe Leu Ala Gln Pro Pro Ser Gly
 165 170 175

20 CAG AGG CGC TAA GGCAGGCTG GGGGGCTTC CTAGGTCATG CCTCTCCCC 636
 Gln Arg Arg *
 180

25 TAGGGAATGG TCCAGCACG AGTGGCAGT TCATTGTGGG GGCTGATG TTGTGCGCTG 696
 GAGGAGGACG GCTTACATGT TTGTTCTGT AGAAAATAAA ACTGAGCTAC GAAAAA 752

30

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

35

- (A) LENGTH: 180 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Met Gln Ala Glu Gly Arg Gly Thr Gly Gly Ser Thr Gly Asp Ala Asp
 1 5 10 15

45 Gly Pro Gly Gly Pro Gly Ile Pro Asp Gly Pro Gly Gly Asn Ala Gly
 20 25 30

Gly Pro Gly Glu Ala Gly Ala Thr Gly Gly Arg Gly Pro Arg Gly Ala
 35 40 45

50 Gly Ala Ala Arg Ala Ser Gly Pro Gly Gly Gly Ala Pro Arg Gly Pro
 50 55 60

55 His Gly Gly Ala Ala Ser Gly Leu Asn Gly Cys Cys Arg Cys Gly Ala
 65 70 75 80

Arg Gly Pro Glu Ser Arg Leu Leu Glu Phe Tyr Leu Ala Met Pro Phe
 85 90 95

11/18

Ala Thr Pro Met Glu Ala Glu Leu Ala Arg Arg Ser Leu Ala Gln Asp
 100 105 110

5 Ala Pro Pro Leu Pro Val Pro Gly Val Leu Leu Lys Glu Phe Thr Val
 115 120 125

Ser Gly Asn Ile Leu Thr Ile Arg Leu Thr Ala Ala Asp His Arg Gln
 130 135 140

10 Leu Gln Leu Ser Ile Ser Ser Cys Leu Gln Gln Leu Ser Leu Leu Met
 145 150 155 160

15 Trp Ile Thr Gln Cys Phe Leu Pro Val Phe Leu Ala Gln Pro Pro Ser
 165 170 175

Gly Gln Arg Arg *

180

20

(2) INFORMATION FOR SEQ ID NO: 9:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 752 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

35

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: homo sapiens

40

(ix) FEATURE:

(A) NAME/KEY: 5'UTR

(B) LOCATION:1..93

45

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION:94..270

50

(ix) FEATURE:

(A) NAME/KEY: 3'UTR

(B) LOCATION:271..752

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

55 ATCTCTGTGG GGGTGAAGT TCTCTGTGAG AGCGGGGAG AGGCTGCGGA GGCATGCGG 60

CGGAGGCGG GGGCAGGG GGTGAGGG GCG ATG CTG ATG GCG CAG GAG GCG 114

Met Leu Met Ala Gln Glu Ala

12/18

		1	5		
5	CTG GCA TTC CTG ATG GGC CAG GGG GCA ATG CTG GCG GGC CAG GAG AGG Leu Ala Phe Leu Met Ala Gln Gly Ala Met Leu Ala Ala Gln Glu Arg	10	15	20	162
10	CGG GTG CCA CGG GCG GCA GAG GTC CCC GGG GCG CAG GGG CAG CAA GGG Arg Val Pro Arg Ala Ala Glu Val Pro Gly Ala Gln Gly Gln Gln Gly	25	30	35	210
15	OCT CGG GGC CGG GAG GAG GCG CCC GCG GGG GTC GCG ATG GCG GCG CGG Pro Arg Gly Arg Glu Glu Ala Pro Arg Gly Val Arg Met Ala Ala Arg	40	45	50	258
20	CTT CAG GGC TGA ATGGATGCTG CAGATGCGGG GCGAGGGGGC CGGAGAGCGG Leu Gln Gly *				310
25	CCGCTGTGAG TTCTACCTCG CCATGCGCTTT CGGACACCC ATGGAGCAG AGCTGGGCGG CAGGAGCGTG GCGCAGGATG CCGCACCGCT TCCCGTGCCA GGGGTGCTTC TGAAGGAGTT				370
30	CACTGTGTCC GGCACATAC TGACTATCCG ACTGACTGCT GCAGACACC GCGAAGTGCA GCTCTCATC AGCTCTGTG TOCAGCAGCT TCCCTGTG ATGAGGATCA CGCAGTGCCT				430
35	TCGCGCGTG TTTTGGCTC AGCTCCCTC AGGGCAGAGG CGCTAAGGCC AGCTGGGCGC CCCTCTCTAG GTCATGCTC CTCCCTAGG GAATGGTCC AGCAGGAGTG GCGAGTTCAT				490
40	TGTTGGGGGC TGATTGTGTG TCGCTGGAGG AGGAGGCGCT ACATGTTTGT TTCTGTAGAA AATAAACTG AGCTACGAAA AA				550
45					610
50					670
55					730
60					752

(2) INFORMATION FOR SEQ ID NO: 10:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 58 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
- 45 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met Leu Met Ala Gln Glu Ala Leu Ala Phe Leu Met Ala Gln Gly Ala
 1 5 10 15
 5 Met Leu Ala Ala Gln Glu Arg Arg Val Pro Arg Ala Ala Glu Val Pro
 20 25 30
 10 Gly Ala Gln Gly Gln Gln Gly Pro Arg Gly Arg Glu Glu Ala Pro Arg
 35 40 45
 Gly Val Arg Met Ala Ala Arg Leu Gln Gly *

15

2) INFORMATION FOR SEQ ID NO: 11:

20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 11 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

30 Met Leu Met Ala Gln Glu Ala Leu Ala Phe Leu
 1 5 10....

35 2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 40 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

45 Leu Met Ala Gln Glu Ala Leu Ala Phe Leu
 1 5 10.

50

(2) INFORMATION FOR SEQ ID NO: 13:

55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

5 GGTGACACTA TAGAGGTAC G 21

10 (2) INFORMATION FOR SEQ ID NO: 14:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

20 TGATGTGCAA CTGAAGCAGG.....20

25 (2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
30 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

35 GCACIGGGTG ATCCACATCA A 21

40 (2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
45 (C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

50 GCATCACA TAGGGAGACA G 21

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: synthetic DNA

GCACATCAGG ATGCGTTTCT CGTGG

25

15

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: synthetic DNA

CACACAAAGC TTGGCTTAGC GGCCTGCGC TG.....32

30

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

- 35 (A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: synthetic DNA

CACACAGGAT CCATGGATGC TGCAGATGG.....30

45

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

- 50 (A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: synthetic DNA

GAAGACATA TGCTGATGGC CCAGGAGGC

29

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

TTAAGATCT CAGAACGGCC CCTGGTCG

28

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

ttactogaga tgctgatggc ccagg.....25

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: synthetic DNA

aaggtacctt gaacggccccc tggctg26

2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

Phe Leu Met Ala Gln Gly Ala Met Leu
1 5 9

5

2) INFORMATION FOR SEQ ID NO: 25:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

20 Ala Met Leu Ala Ala Gln Glu Arg Arg Val
1 5 10

25

2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acids

30

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Met Leu Ala Ala Gln Glu Arg Arg Val
1 5 9

40

2) INFORMATION FOR SEQ ID NO: 27:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

55 Tyr Tyr Met Asn Gly Thr Met Ser Gln Val
1 5 10

2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Glu Val Asp Pro Ile Gly His Leu Tyr
1 5 9

15

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 99/07832

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WANG, R.-F. ET AL.: "A breast and melanoma-shared tumor antigen: T cell responses to antigenic peptides translated from different open reading frames." JOURNAL OF IMMUNOLOGY, vol. 161, 1 October 1998 (1998-10-01), pages 3596-606, XP002096029 the whole document	12-14
X	WO 98 32855 A (GODELAINE DANIELE ;LETHE BERNARD (BE); LUCAS SOPHIE (BE); SMET CHA) 30 July 1998 (1998-07-30) see whole document, particularly the claims	12-14



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 March 2000

Date of mailing of the international search report

24/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Smalt, R

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 99/07832

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VAN ELSAS, A. ET AL.: "Transfection of IL-2 augments CTL response to human melanoma cells in vitro: immunological characterization of a melanoma vaccine." JOURNAL OF IMMUNOTHERAPY, vol. 20, no. 5, September 1997 (1997-09), pages 343-53, XP002096030 cited in the application abstract; figures 6B,7	
P,X	AARNOUDSE, C. A. ET AL.: "Interleukin-2-induced, melanoma-specific T cells recognize CAMEL, an unexpected translation product of LAGE-1." INTERNATIONAL JOURNAL OF CANCER, vol. 82, no. 3, 30 July 1999 (1999-07-30), pages 442-8, XP000867744 the whole document	1-4,9-14
P,X	WO 99 18206 A (WANG RONG FU ;GOVERNMENT OF THE UNITED STATE (US); ROSENBERG STEVE) 15 April 1999 (1999-04-15) claims 24,26,27,37; figures 3A,7A	3,9,11

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 99/07832

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0 9832855 A	30-07-1998	US 5811519 A	22-09-1998
		AU 6042198 A	18-08-1998
		EP 0970206 A	12-01-2000
		ZA 9800656 A	17-08-1998
W0 9918206 A	15-04-1999	AU 9572098 A	27-04-1999